

On refined enumerations of totally symmetric self-complementary plane partitions II

Masao Ishikawa

Faculty of Education, Tottori University
Koyama, Tottori, Japan
ishikawa@fed.tottori-u.ac.jp

Mathematics Subject Classifications: Primary 05A15; Secondary 05A17, 05E05, 05E10.

Keywords: totally symmetric self-complementary plane partitions; determinantal formulae; half-turn (or vertically) symmetric alternating sign matrices; twisted Bender-Knuth involution; domino plane partitions.

Abstract

In this paper we settle a weak version of a conjecture (i.e. Conjecture 6) by Mills, Robbins and Rumsey in the paper “Self-complementary totally symmetric plane partitions” *J. Combin. Theory Ser. A* **42**, 277–292. In other words we show that the number of shifted plane partitions invariant under the involution γ is equal to the number of alternating sign matrices invariant under the vertical flip. We also give a determinant expression of the general conjecture (Conjecture 6), but this determinant is still hard to evaluate. In this paper we introduce two new classes of domino plane partitions, one has the same cardinality as the set of half-turn symmetric alternating sign matrices and the other has the same cardinality as the set of vertically symmetric alternating sign matrices.

Contents

1	Introduction	1
2	Preliminaries	4
3	Bijections and Statistics	6
4	A twisted Bender-Knuth involution	8
5	Twisted domino plane partitions	13
6	Domino plane partitions	15
7	Determinantal formulae	18

1 Introduction

A totally symmetric self-complementary plane partition is, by definition, a plane partition which is invariant under permutation of the three axes and which is equal

to its complement (cf. [6, 8, 17, 18, 22, 23]). The number of totally symmetric self-complementary plane partitions is known to be the same as that for alternating sign matrices and descending plane partitions. But, there are still several interesting conjectures concerning totally symmetric self-complementary plane partitions (see [17]). This paper is the succession of my previous paper [8] in which we obtain Pfaffian formulae and constant term identities for Conjecture 2, Conjecture 3 and Conjecture 7 as an application of the minor summation formulas of Pfaffians obtained in [9, 10]. In this paper we are mainly concerned with two other conjectures, i.e. Conjecture 4 and Conjecture 6, by Mills, Robbins and Rumsey in the paper [17]. We will obtain a determinantal formula for Conjecture 6 as an application of essentially a Binet-Cauchy type formula. We also introduce two new classes of domino plane partitions which seemingly look closely related to Conjecture 4 and Conjecture 6.

In [17] Mills, Robbins and Rumsey have introduced a set of triangular shifted plane partitions, which is bijective to the set of totally symmetric self-complementary plane partitions. In this paper we denote this set by \mathcal{B}_n , which is defined to be the set of triangular shifted plane partitions $b = (b_{ij})_{1 \leq i \leq j \leq n-1}$ whose parts are $\leq n$, weakly decreasing along rows and columns, and all parts in row i are $\geq n - i$. For $b = (b_{ij})_{1 \leq i \leq j \leq n-1}$ in \mathcal{B}_n and $k \geq 1$, let

$$U_k(b) = \sum_{t=1}^{n-k} (b_{t,t+k-1} - b_{t,t+k}) + \sum_{t=n-k+1}^{n-1} \chi\{b_{t,n-1} > n - t\}. \quad (1.1)$$

Here we use the convention that $b_{i,n} = n - i$ for all i and $b_{0,j} = n$ for all j . They have also introduced two involutions ρ and γ of \mathcal{B}_n onto itself, and conjectured that they correspond to the half turn and the vertical flip of the alternating sign matrices. These involutions are defined as follows. Let $b = (b_{ij})_{1 \leq i \leq j \leq n-1}$ be an element of \mathcal{B}_n and let b_{ij} be a part of b off the main diagonal. Then the *flip* of the part b_{ij} is the operation of replacing b_{ij} by b'_{ij} where

$$b'_{ij} + b_{ij} = \min(b_{i-1,j}, b_{i,j-1}) + \max(b_{i,j+1}, b_{i+1,j}). \quad (1.2)$$

When the part is in the main diagonal, the flip of a part b_{ii} is the operation replacing b_{ii} by b'_{ii} where

$$b'_{ii} + b_{ii} = b_{i-1,i} + b_{i,i+1}. \quad (1.3)$$

An operation π_r is defined to be a map $\mathcal{B}_n \rightarrow \mathcal{B}_n$ where $\pi_r(b)$ is the result of flipping all the $b_{i,i+r-1}$, $1 \leq i \leq n - r$. We can introduce two involutions ρ and γ as follows:

$$\rho = \pi_2 \pi_4 \pi_6 \cdots \quad (1.4)$$

$$\gamma = \pi_1 \pi_3 \pi_5 \cdots \quad (1.5)$$

(see ([17, pp.284,286])). Mills, Robbins and Rumsey conjectured that the invariants of ρ in \mathcal{B}_n correspond to the half-turn symmetric alternating sign matrices, and the invariants of γ in \mathcal{B}_n correspond to the vertically symmetric alternating sign matrices. Let \mathcal{B}_n^ρ (resp. \mathcal{B}_n^γ) denotes the set of elements in \mathcal{B}_n invariant under ρ (resp. γ). These conjectures are stated as follows. Here, for the definition of the numbers A_n^{HTS} , A_{2n+1}^{VS} and the polynomials $A_n^{\text{HTS}}(t)$, $A_{2n+1}^{\text{VS}}(t)$, see the next section.

Conjecture 1.1. ([17, pp.285, Conjecture 4]) Let $n \geq 2$ and r , $0 \leq r < n$, be integers. Then the number of elements of \mathcal{B}_n with $\rho(b) = b$ and $U_1(b) = r$ would be the same as the number of $n \times n$ alternating sign matrices invariant under the half turn and satisfying $a_{1,r+1} = 1$. Namely, $\sum_{b \in \mathcal{B}_n^\rho} t^{U_1(b)} = A_n^{\text{HTS}}(t)$ would hold.

Conjecture 1.2. ([17, pp.286, Conjecture 6]) Let $n \geq 1$ be an integer and r , $1 \leq r \leq 2n - 1$, be an integer. Then the number of elements of \mathcal{B}_{2n+1} with $\gamma(b) = b$ and $U_2(b) = r - 1$ would be the same as the number of $n \times n$ alternating sign matrices with $a_{i1} = 1$ and invariant under the vertical flip. Namely, $\sum_{b \in \mathcal{B}_{2n+1}^\gamma} t^{U_2(b)} = A_{2n+1}^{\text{VS}}(t)$ would hold.

In [8] we have introduced a set \mathcal{P}_n of column strict plane partitions which is bijective with the set \mathcal{B}_n of triangular symmetric plane partitions (Theorem 3.3). Namely \mathcal{P}_n is, by definition, the set of column-strict plane partitions in which each entry in the j th column does not exceed $n - j$. We recall these plane partitions, the bijections and the statistics in Section 3. In Section 4, we translate the involution π_r in the words of \mathcal{P}_n , and find that the involution π_r correspond to a Bender-Knuth type involution $\tilde{\pi}_r$ which swaps i and $i - 1$ in a column-strict plane partition in \mathcal{P}_n . Let

$$\tilde{\rho} = \tilde{\pi}_2 \tilde{\pi}_4 \tilde{\pi}_6 \cdots, \quad (1.6)$$

$$\tilde{\gamma} = \tilde{\pi}_1 \tilde{\pi}_3 \tilde{\pi}_5 \cdots, \quad (1.7)$$

and let $\mathcal{P}_n^{\tilde{\rho}}$ (resp. $\mathcal{P}_n^{\tilde{\gamma}}$) denotes the set of invariants of $\tilde{\rho}$ (resp. $\tilde{\gamma}$). Since this Bender-Knuth type involution $\tilde{\pi}_r$ is, in a sense, “twisted” (a little different from the ordinary one), we will see that the set $\mathcal{P}_n^{\tilde{\rho}}$ is naturally bijective to a set \mathcal{G}_n of “twisted” domino plane partitions in Section 5 (see Theorem 5.2). In Section 6, we introduce two new classes of domino plane partitions, i.e. \mathcal{D}_n^{R} and \mathcal{D}_n^{C} . Namely, \mathcal{D}_n^{R} (resp. \mathcal{D}_n^{C}) is defined to be the set of column-strict domino plane partitions whose entries in the j th column are $\leq \lceil (n - j)/2 \rceil$ and with all rows (resp. columns) of even length. We also construct a natural bijection between $\mathcal{P}_{2n+1}^{\tilde{\gamma}}$ and $\mathcal{D}_{2n-1}^{\text{R}}$ (see Theorem 6.2). It seems that \mathcal{G}_n and \mathcal{D}_n^{C} ($n \geq 1$) have the same number of elements from examples, but we don’t know how to construct a bijection between them at this point (see Conjecture 6.3). The following diagram of the bijections give a one-to-one correspondence between the set $\mathcal{B}_{2n+1}^\gamma$ of triangular shifted plane partitions invariant under γ and the set $\mathcal{D}_{2n-1}^{\text{R}}$ of domino plane partitions:

$$\mathcal{B}_{2n+1}^\gamma \xrightarrow{\varphi_{2n+1}} \mathcal{P}_{2n+1}^{\tilde{\gamma}} \xrightarrow{\tau_{2n+1}} \mathcal{D}_{2n-1}^{\text{R}} \xrightarrow{\Phi} \mathcal{Q}_{2n-1}^{\text{H}}$$

Meanwhile, we obtain the following one-to-one correspondence

$$\mathcal{B}_n^\rho \xrightarrow{\varphi_n} \mathcal{P}_n^{\tilde{\rho}} \xrightarrow{\psi_n} \mathcal{G}_n \dashrightarrow \mathcal{D}_n^{\text{C}} \xrightarrow{\Phi} \mathcal{Q}_n^{\text{V}}$$

between the set \mathcal{B}_n^ρ of triangular shifted plane partitions invariant under ρ and the set \mathcal{G}_n of “twisted” domino plane partitions, whereas we don’t know the missing bijection between \mathcal{G}_n and \mathcal{D}_n^{C} . In Section 7, we use a plane partition analogue Φ of the Stanton-White bijection which maps a domino plane partition to a paired plane partition (see [7, 25]), that enable us to define a bijection between \mathcal{D}_n^{R} (resp.

\mathcal{D}_n^C) of domino plane partitions and \mathcal{Q}_n^H (resp. \mathcal{Q}_n^V) of paired plane partitions. (see Theorem 7.2). Using the generating functions obtained in [8], we obtain determinantal formulae for the generating functions of these sets of paired plane partitions (see Corollary 7.7). As a special case we show that Conjecture 1.2 reduce to the evaluation of the determinant in the following theorem:

Theorem 1.3. Let $n \geq 2$ be a positive integer. Let $\det R_n^o(t) = (R_{i,j}^o)_{0 \leq i,j \leq n}$ be the $n \times n$ matrix where

$$R_{i,j}^o = \binom{i+j-1}{2i-j} + \left\{ \binom{i+j-1}{2i-j-1} + \binom{i+j-1}{2i-j+1} \right\} t + \binom{i+j-1}{2i-j} t^2$$

with the convention that $R_{0,0}^o = R_{0,1}^o = 1$. Then we obtain

$$\sum_{b \in \mathcal{B}_{2n+1}^*} t^{U_2(b)} = \det R_n^o(t). \quad (1.8)$$

(See Corollary 7.8(ii)).

Thus Conjecture 1.2 reduce to prove that $\det D_n(t) = A_{2n+1}^{\text{VS}}(t)$ would hold. We also obtain a similar formula for the generating function of \mathcal{Q}_n^V (see Corollary 7.8). This determinant is also conjectured to be $A_n^{\text{HTS}}(t)$, but still hard to evaluate (see Conjecture 7.9). (About determinant evaluation the reader can consult [12]). Meanwhile, when $t = 1$, we will find that we can reduce the evaluation of these determinants to the Andrews-Burge determinant (7.21) (see [1, 2, 4, 18]) and we obtain the result that the number of elements in \mathcal{Q}_n^R (resp. \mathcal{Q}_n^C) is equal to the number A_n^{VS} (resp. A_n^{HTS}) of vertically (resp. half-turn) symmetric alternating sign matrices. Thus we prove Conjecture 1.2 is true when $t = 1$, whereas we can't say it for Conjecture 1.1 because of the missing bijection. Anyway we define a new object \mathcal{D}_n^C of domino plane partitions which has the same cardinality with the set of half-turn symmetric alternating sign matrices. In the study of these several classes of plane partitions, we will see they possess many similarities with Young tableaux and Schur functions, but sometimes they are twisted and have mysterious coincidences which we can't explain.

2 Preliminaries

Let $A_n(t)$ be the polynomial defined by

$$A_n(t) = \frac{A_n}{\binom{3n-2}{n-1}} \sum_{r=1}^n \binom{n+r-2}{n-1} \binom{2n-1-r}{n-1} t^{r-1}, \quad (2.1)$$

where A_n is the number defined by $A_n = \prod_{i=0}^{n-1} \frac{(3i+1)!}{(n+i)!}$. It is well-known that A_n is the number of alternating sign matrices and $A_n(t)$ is the refined ASM distribution (see [13, 16, 22, 27]). Let A_{2n}^{HTS} be the number defined by

$$A_{2n}^{\text{HTS}} = \prod_{i=0}^{n-1} \frac{(3i)!(3i+2)!}{\{(n+i)!\}^2} \quad \text{and} \quad A_{2n+1}^{\text{HTS}} = \frac{n!(3n)!}{\{(2n)!\}^2} \cdot A_{2n}^{\text{HTS}}, \quad (2.2)$$

which is known to be the number of half-turn symmetric alternating sign matrices (see [14, 19, 21, 26]). The first few terms of (2.2) are 1, 2, 3, 10, 25, 140, 588. We follow [21] and define the polynomial $\tilde{A}_n^{\text{HTS}}(t)$ by

$$\frac{\tilde{A}_{2n}^{\text{HTS}}(t)}{\tilde{A}_{2n}^{\text{HTS}}} = \frac{(3n-2)(2n-1)!}{(n-1)!(3n-1)!} \sum_{r=0}^n \frac{\{n(n-1) - nr + r^2\}(n+r-2)!(2n-r-2)!}{r!(n-r)!} t^r \quad (2.3)$$

where $\tilde{A}_{2n}^{\text{HTS}} = \prod_{i=0}^{n-1} \frac{(3i)!(3i+2)!}{(3i+1)!(n+i)!}$. Let

$$A_{2n}^{\text{HTS}}(t) = \tilde{A}_{2n}^{\text{HTS}}(t)A_n(t), \quad (2.4)$$

and

$$A_{2n+1}^{\text{HTS}}(t) = \frac{1}{3} \left\{ A_{n+1}(t)\tilde{A}_{2n}^{\text{HTS}}(t) + A_n(t)\tilde{A}_{2n+2}^{\text{HTS}}(t) \right\}, \quad (2.5)$$

which is known to be the refined enumeration of half-turn symmetric alternating sign matrices weighted by the distribution of one in the top row. The first few terms of (2.4) and (2.5) are $A_2^{\text{HTS}}(t) = 1+t$, $A_3^{\text{HTS}}(t) = 1+t+t^2$, $A_4^{\text{HTS}}(t) = 2+3t+3t^2+2t^3$, $A_5^{\text{HTS}}(t) = 3+6t+7t^2+6t^3+3t^4$.

We follow [20] and define A_{2n+1}^{VS} and $A_{2n+1}^{\text{VS}}(t)$ as follows. Let A_{2n+1}^{VS} be the number given by

$$A_{2n+1}^{\text{VS}} = (-3)^{n^2} \prod_{\substack{1 \leq i, j \leq 2n+1 \\ 2 \mid j}} \frac{3(j-i)+1}{j-i+2n+1} = \frac{1}{2^n} \prod_{k=1}^n \frac{(6k-2)!(2k-1)!}{(4k-2)!(4k-1)!}. \quad (2.6)$$

This number A_{2n+1}^{VS} is equal to the number of vertically symmetric alternating sign matrices of size $2n+1$ (see [14, 19, 20]), and the first few terms of (2.6) are 1, 3, 26, 646 and 45885. Let $A_{2n+1}^{\text{VS}}(t)$ be the polynomial

$$A_{2n+1}^{\text{VS}}(t) = \frac{A_{2n-1}^{\text{VS}}}{(4n-2)!} \sum_{r=1}^{2n} t^{r-1} \sum_{k=1}^r (-1)^{r+k} \frac{(2n+k-2)!(4n-k-1)!}{(k-1)!(2n-k)!}, \quad (2.7)$$

which is known to be the refined enumeration of vertically symmetric alternating sign matrices weighted by the distribution of one in the first column (see [20]). For instance, the first few terms of (2.7) are $A_3^{\text{VS}}(t) = 1$, $A_5^{\text{VS}}(t) = 1+t+t^2$, $A_7^{\text{VS}}(t) = 3+6t+8t^2+6t^3+3t^4$ and $A_9^{\text{VS}}(t) = 26+78t+138t^2+162t^3+138t^4+78t^5+26t^6$, and we have $A_{2n+1}^{\text{VS}}(1) = A_{2n+1}^{\text{VS}}$.

Next we recall the terminology of partitions and plane partitions. We follow the notation in Macdonald [15] and Stanley [24]. If the reader is familiar with the notion, he can skip the rest of the section. Let \mathbb{P} denote the set of positive integers. A partition is a sequence $\lambda = (\lambda_1, \lambda_2, \dots)$ of non-negative integers in non-increasing order: $\lambda_1 \geq \lambda_2 \geq \dots$ and containing only finitely many non-zero terms. The non-zero λ_i are called the *parts* of λ . The number of parts is the *length* of λ , denoted by $\ell(\lambda)$; and the sum of parts is the *weight* of λ , denoted by $|\lambda|$. The *diagram* of a partition λ may be formally defined as the set of lattice points $(i, j) \in \mathbb{P}^2$ such that $1 \leq j \leq \lambda_i$. We identify λ with its diagram. The *conjugate* of a partition λ is the partition λ' whose diagram is the transpose of the diagram of λ . A partition with

distinct parts is called a *strict partition*. The *shifted diagram* of a strict partition μ is the set of lattice points $(i, j) \in \mathbb{P}^2$ such that $i \leq j \leq \mu_i + i$. We identify a strict partition with its shifted diagram.

A *plane partition* is an array $\pi = (\pi_{ij})_{i,j \geq 1}$ of nonnegative integers such that π has finite support (i.e. finitely many nonzero entries) and is weakly decreasing in rows and columns. If $\sum_{i,j \geq 1} \pi_{ij} = n$, then we write $|\pi| = n$ and say that π is a plane partition of n , or π has *weight* n . A *part* of a plane partition $\pi = (\pi_{ij})_{i,j \geq 1}$ is a positive entry $\pi_{ij} > 0$. The *shape* of π is the ordinary partition λ for which π has λ_i nonzero parts in the i th row. We denote the shape of π by $\text{sh}(\pi)$. We also say that π has r rows if $r = \ell(\lambda)$. Similarly, π has s columns if $s = \ell(\lambda')$. A plane partition is said to be *column-strict* if it is strictly decreasing in columns.

Let μ be a strict partition. A *shifted plane partition* τ of *shifted shape* μ is an arbitrary filling of the cells of μ with nonnegative integers such that each entry is weakly decreasing in rows and columns. In this paper we allow parts to be zero for shifted plane partitions of a fixed shifted shape μ .

3 Bijections and Statistics

First we recall the results we obtained in the preceding paper [8]. We defined the set $\mathcal{P}_{n,m}$ of plane partitions and studied it intensively. This set $\mathcal{P}_{n,m}$ is also the main object we study in this paper:

Definition 3.1. Let m and $n \geq 1$ be nonnegative integers. Let $\mathcal{P}_{n,m}$ denote the set of column-strict plane partitions $c = (c_{ij})_{1 \leq i,j}$ subject to the constraints that

(C1) c has at most n columns;

(C2) each part in the j th column of c does not exceed $n + m - j$.

An element of $\mathcal{P}_{n,m}$ is called a *restricted column-strict plane partition*. When $m = 0$, we write \mathcal{P}_n for $\mathcal{P}_{n,0}$. If a part in the j th column of c is equal to $n + m - j$, we call the part a *saturated part*.

Let $c = (c_{ij})_{1 \leq i \leq n+m, 1 \leq j \leq n}$ be a plane partition in $\mathcal{P}_{n,m}$ and let k be a positive integer. Let $c_{\geq k}$ denote the plane partition formed by the parts $\geq k$. Let

$$\theta_i(c_{\geq k}) = \#\{l : c_{i,l} \geq k\} \quad (3.1)$$

denote the length of the i th row of $c_{\geq k}$, i.e. the rightmost column containing a letter $\geq k$ in the i th row of c .

Let r be an integer such that $1 \leq r \leq n + m$. For $c \in \mathcal{P}_{n,m}$ let $\bar{U}_r(c)$ be the number of parts equal to r plus the number of saturated parts less than r , i.e.

$$\bar{U}_r(c) = \#\{(i, j) : c_{ij} = r\} + \#\{1 \leq k < r : c_{1,n+m-k} = k\}. \quad (3.2)$$

Especially $\bar{U}_1(c)$ is the number of 1's in c and $\bar{U}_{n+m}(c)$ is the number of saturated

parts in c . For example, let

6	6	4	4	3	1	1
5	3	3	2	1		
3	2	2	1			
1	1					

be an element c of \mathcal{P}_8 , then, the bold faced entries are the saturated parts. Thus we have $\bar{U}_1(c) = 6$, $\bar{U}_2(c) = \bar{U}_4(c) = \bar{U}_5(c) = \bar{U}_7(c) = \bar{U}_8(c) = 4$ and $\bar{U}_3(c) = \bar{U}_6(c) = 5$.

We also defined the following set $\mathcal{B}_{n,m}$ of shifted plane partitions in [8], which is a generalization of \mathcal{B}_n defined in [17, pp.281].

Definition 3.2. (See [11, Theorem 1]). Let m and $n \geq 1$ be nonnegative integers. Let $\mathcal{B}_{n,m}$ denote the set of shifted plane partitions $b = (b_{ij})_{1 \leq i \leq j}$ subject to the constraints that

(B1) the shifted shape of b is $(n + m - 1, n + m - 2, \dots, 2, 1)$;

(B2) $\max\{n - i, 0\} \leq b_{ij} \leq n$ for $1 \leq i \leq j \leq n + m - 1$.

When $m = 0$, we write \mathcal{B}_n for $\mathcal{B}_{n,0}$. In this paper we call an element of $\mathcal{B}_{n,m}$ a *triangular shifted plane partition* (abbreviated to TSPP).

We use the convention that $b_{i,n+m} = n - i$ for all i and $b_{0,j} = n$ for all j . For a $b = (b_{ij})_{1 \leq i \leq j \leq n+m-1}$ in $\mathcal{B}_{n,m}$ and an integer r such that $1 \leq r \leq n + m$, let

$$U_r(b) = \sum_{t=1}^{n+m-r} (b_{t,t+r-1} - b_{t,t+r}) + \sum_{t=n+m-r+1}^{n+m-1} \chi\{b_{t,n+m-1} > n - t\}. \quad (3.3)$$

We put $\bar{U}_r(b) = n + m - 1 - U_r(b)$. For example, let

8	8	8	8	8	8	8
	8	8	8	8	7	6
		8	8	7	7	6
			6	5	5	4
				4	4	3
					3	3
						1

be an element b of \mathcal{B}_8 , then we have $U_1(b) = 1$, $U_2(b) = U_4(b) = U_5(b) = U_7(b) = U_8(b) = 3$ and $U_3(b) = U_6(b) = 2$.

In [8] we have established a bijection between $\mathcal{B}_{n,m}$ and $\mathcal{P}_{n,m}$, and proved that these statistics agree.

Theorem 3.3. Let m and $n \geq 1$ be nonnegative integers and let $c = (c_{ij})_{1 \leq i \leq n+m, 1 \leq j \leq n}$ be a RCSPP in $\mathcal{P}_{n,m}$. Associate to the array $c = (c_{ij})_{1 \leq i \leq n+m, 1 \leq j \leq n}$ the array $b = (b_{ij})_{1 \leq i \leq j \leq n+m-1}$ defined by

$$n - b_{ij} = \theta_{n+m-j}(c_{\geq 1-i+j}) \quad (3.4)$$

with $1 \leq i \leq j \leq n + m - 1$. Then b is in $\mathcal{B}_{n,m}$, and this mapping $\varphi_{n,m}$, which associate to a RCSPP c the TSPP $b = \varphi_{n,m}(c)$, is a bijection of $\mathcal{P}_{n,m}$ onto $\mathcal{B}_{n,m}$. Further, by this bijection, we have $\overline{U}_r(\varphi_{n,m}(c)) = \overline{U}_r(c)$ for any $c \in \mathcal{P}_{n,m}$.

For instance, the b and c in the above examples correspond to each other by this bijection between \mathcal{B}_8 and \mathcal{P}_8 . In [8, Section 2] we defined a set $\mathcal{T}_{n,m}$ of totally symmetric plane partitions and constructed the bijections $\mathcal{T}_{n,m} \leftrightarrow \mathcal{B}_{n,m}$ and $\mathcal{T}_{n,m} \leftrightarrow \mathcal{P}_{n,m}$. Thus the study of totally symmetric plane partitions reduce to the study of restricted column-strict plane partitions.

4 A twisted Bender-Knuth involution

A classical method to prove that a Schur function is symmetric is to define involutions s_i on tableaux which swaps the number of i 's and $(i - 1)$'s, for each i . This is well-known as the Bender-Knuth involution ([5]). In this section we define a twisted Bender-Knuth involution $\tilde{\pi}_r$ of the set $\mathcal{P}_{n,m}$ of RCSPPs and show that it correspond to the involution π_r of $\mathcal{B}_{n,m}$. We continue to use the convention that $b_{i,n+m} = n - i$ for all i and $b_{0,j} = n$ for all j .

In [17], Mills, Robbins and Rumsey have introduced the notion of flip for \mathcal{B}_n . We can naturally generalize this notion to $\mathcal{B}_{n,m}$ by the same equations (1.2) and (1.3), whereas we have to be careful about the range of m . Let m and $n \geq 1$ be non-negative integers. Let $b = (b_{ij})_{1 \leq i \leq j \leq n+m-1}$ be an element of $\mathcal{B}_{n,m}$ and let $1 \leq i < j \leq n + m - 1$ so that b_{ij} is a part of b off the main diagonal. We define the *flip* of the part b_{ij} as the operation of replacing b_{ij} with b'_{ij} with (1.2). Note that this operation is always well-defined since b'_{ij} satisfies the axiom (B2), and the result of flipping b_{ij} is a shifted plane partition. In fact $b \in \mathcal{B}_{n,m}$ implies that $\min(b_{i-1,j}, b_{i,j-1}) \geq b_{ij} \geq \max(b_{i,j+1}, b_{i+1,j})$ so that we have $n \geq \min(b_{i-1,j}, b_{i,j-1}) \geq b'_{ij} \geq \max(b_{i,j+1}, b_{i+1,j}) \geq 0$ and $b'_{ij} \geq b_{i,j+1} \geq n - i$.

When the part is in the main diagonal, we define the flip of a part b_{ii} as the operation replacing b_{ii} with b'_{ii} with (1.3). Note that the result of flipping a part b_{ii} in the main diagonal may violate the axiom (B2) unless $m = 0$ or $m = 1$. In fact, if $m \geq 2$, then $b'_{n+m-1,n+m-1}$ can be negative since $b_{n+m-1,n+m} = 1 - m < 0$. Thus, hereafter, we assume $m = 0$ or 1 when we consider a flip of a part in the main diagonal.

Let $1 \leq r \leq n + m$ and $b = (b_{ij})_{1 \leq i \leq j \leq n+m-1} \in \mathcal{B}_{n,m}$. Define an operation $\pi_r : \mathcal{B}_{n,m} \rightarrow \mathcal{B}_{n,m}$ by $b \mapsto \pi_r(b)$ where $\pi_r(b)$ is the result of flipping all the $b_{i,i+r-1}$, $1 \leq i \leq n + m - r$. Since none of these parts of b are neighbors, the result is independent of the order in which the flips are applied, and this operation π_r is evidently an involution, i.e. $\pi_r^2 = id$. For instance, the following TSPP $b \in \mathcal{B}_6$ is mapped to the following TSPP $\pi_2(b)$ by the involution π_2 , and to the following TSPP $\pi_1(b)$ by the involution π_1 ;

$$\begin{array}{rcc}
 \begin{array}{cccccc}
 6 & 6 & 6 & 6 & 5 & \\
 & 6 & 5 & 5 & 5 & \\
 & & 4 & 4 & 4 & , \\
 & & & 4 & 4 & \\
 & & & & 1 &
 \end{array} & , & \pi_2(b) = \begin{array}{cccccc}
 6 & \mathbf{6} & 6 & 6 & 5 & \\
 & 6 & \mathbf{6} & 5 & 5 & \\
 & & 4 & \mathbf{4} & 4 & , \\
 & & & 4 & \mathbf{2} & \\
 & & & & 1 &
 \end{array} & , & \pi_1(b) = \begin{array}{cccccc}
 \mathbf{6} & 6 & 6 & 6 & 5 & \\
 & \mathbf{5} & 6 & 5 & 5 & \\
 & & \mathbf{5} & 4 & 4 & . \\
 & & & \mathbf{4} & 2 & \\
 & & & & \mathbf{4} &
 \end{array}
 \end{array}$$

Now we define a Bender-Knuth type involution $\tilde{\pi}_r : \mathcal{P}_{n,m} \rightarrow \mathcal{P}_{n,m}$. This involution $\tilde{\pi}_r$ is an “almost Bender-Knuth involution” which swaps r ’s and $r - 1$ ’s except the fact that it does not count a saturated $r - 1$. In fact, if it did convert a saturated $r - 1$ of c in $\mathcal{P}_{n,m}$, the resulting plane partition could violate the axiom of $\mathcal{P}_{n,m}$. Let see the exact definition. Let $2 \leq r \leq n + m$ and $c \in \mathcal{P}_{n,m}$. Consider the parts of c equal to r or $r - 1$. Since c is column-strict, some columns of c will contain neither r nor $r - 1$, while some others will contain one r and one $r - 1$. These columns we ignore. We also ignore an $r - 1$ in column $n + m - r + 1$, i.e. we ignore a saturated part which is equal to $r - 1$ because a saturated $r - 1$ can’t be changed to r . The remaining parts equal to r or $r - 1$ occur once in each column. Assume row i has a certain number k of r ’s followed by a certain number l of $r - 1$ ’s. Note that we don’t count an $r - 1$ if it is saturated so that a saturated $r - 1$ always remains untouched. For example, the three consecutive rows $i - 1$, i and $i + 1$ of c could look as follows. In row i , convert the k r ’s and l $r - 1$ ’s to l r ’s and k $r - 1$ ’s. It is easy to see that

$$\begin{array}{c}
 i - 1 \\
 i \\
 i + 1
 \end{array}
 \left\| \begin{array}{ccc}
 \vdots & & \vdots \\
 r & \dots & r \\
 r - 1 & \dots & r - 1
 \end{array} \right.
 r \dots r \quad r - 1 \dots r - 1
 \begin{array}{|c|c|c|}
 \hline
 r & \dots & r \\
 \hline
 r - 1 & \dots & r - 1 \\
 \hline
 \end{array}$$

the resulting array satisfies the axioms (C1) and (C2) of $\mathcal{P}_{n,m}$. Define an operation $\tilde{\pi}_r : \mathcal{P}_{n,m} \rightarrow \mathcal{P}_{n,m}$ by $c \mapsto \tilde{\pi}_r(c)$ where $\tilde{\pi}_r(c)$ is the result of swapping r ’s and $r - 1$ ’s in row i of c by this twisted rule for $1 \leq i \leq n + m - r$. For example, if $n = 6$, $m = 0$ and $r = 2$, then the left below RCSPP c corresponds to the right below RCSPP $\tilde{\pi}_2(c)$ by $\tilde{\pi}_2$.

$$c = \begin{array}{|c|c|c|c|c|}
 \hline
 \mathbf{5} & 3 & 1 & 1 & \mathbf{1} \\
 \hline
 3 & 2 & & & \\
 \hline
 2 & 1 & & & \\
 \hline
 \end{array}
 \quad
 \tilde{\pi}_2(c) = \begin{array}{|c|c|c|c|c|}
 \hline
 \mathbf{5} & 3 & 2 & \mathbf{2} & \mathbf{1} \\
 \hline
 3 & 2 & & & \\
 \hline
 1 & 1 & & & \\
 \hline
 \end{array}$$

Next assume $m = 0$ or 1 , and let c be a plane partition in $\mathcal{P}_{n,m}$. Set $\lambda_i = \theta_i(c_{\geq 2})$ to be the number of parts ≥ 2 in the i th row of c . Assume the i th row contains a certain number k of 1 ’s followed by a certain number l of blank positions which 1 ’s can be put in, so that we have $k + l = n + m - 1 - \lambda_1$ if $i = 1$, $k + l = \lambda_{i-1} - \lambda_i$ otherwise. Change the number of 1 ’s from k to l in row i for $1 \leq i \leq n + m - 1$. It is also easy to see that the resulting array, say $\tilde{\pi}_1(c)$, satisfies the constraints (C1) and (C2). For example, if c is as above, then $\tilde{\pi}_1(c)$ is as follows:

$$\tilde{\pi}_1(c) = \begin{array}{|c|c|}
 \hline
 \mathbf{5} & 3 \\
 \hline
 3 & 2 \\
 \hline
 2 & \\
 \hline
 1 & \\
 \hline
 \end{array}
 .$$

This mapping $\tilde{\pi}_1 : \mathcal{P}_{n,m} \rightarrow \mathcal{P}_{n,m}$ is well-defined for $m = 0, 1$, and is evidently an involution. We call this involution $\tilde{\pi}_r$, $1 \leq r \leq n + m$, a *twisted Bender-Knuth*

involution (abbreviated to the *TBK involution*). Note that when $r = n + m$, π_{n+m} and $\tilde{\pi}_{n+m}$ are both the identity mapping since there are no parts affected by the operations. The following proposition corresponds to Theorem 2 of [17, pp.283], whereas there is no need of proof since it is clear from the above definition.

Proposition 4.1. Let m and $n \geq 1$ be non-negative integers. Let $2 \leq r \leq n + m$ and let c in $\mathcal{P}_{n,m}$. Then

$$\overline{U}_r(\tilde{\pi}_r(c)) = \overline{U}_{r-1}(c) \text{ and } \overline{U}_r(c) = \overline{U}_{r-1}(\tilde{\pi}_r(c)) \quad \square$$

The following theorem tells us that the involution π_r of $\mathcal{B}_{n,m}$ corresponds to $\tilde{\pi}_r$ of $\mathcal{P}_{n,m}$ if we identify $\mathcal{B}_{n,m}$ with $\mathcal{P}_{n,m}$ by the bijection $\varphi_{n,m}$ defined in Corollary 3.3.

Theorem 4.2. Let m and $n \geq 1$ be non-negative integers and let $1 \leq r \leq n + m$. Assume $m = 0$ or 1 if $r = 1$. Then we have

$$\pi_r(\varphi_{n,m}(c)) = \varphi_{n,m}(\tilde{\pi}_r(c)).$$

In [17, pp.284], Mills, Robbins and Rumsey defined an involution ρ of \mathcal{B}_n by $\rho = \pi_2\pi_4\cdots$ and presented a conjecture (Conjecture 1.1) that this involution ρ corresponds to the half turn of an alternating matrix. We naturally generalize this definition to $\mathcal{B}_{n,m}$ and use the same symbol for the involution $\rho : \mathcal{B}_{n,m} \rightarrow \mathcal{B}_{n,m}$ defined by

$$\rho = \pi_2\pi_4\cdots \tag{4.1}$$

where the product is over all π_i with i even and $\leq n$. Let $\mathcal{B}_{n,m}^\rho$ denote the set of elements of $\mathcal{B}_{n,m}$ invariant under ρ , i.e. $\mathcal{B}_{n,m}^\rho = \{b \in \mathcal{B}_{n,m} \mid \rho(b) = b\}$. For instance, the following TSPP is an element of \mathcal{B}_8^ρ :

$$\begin{array}{cccccc} 8 & \mathbf{8} & 8 & \mathbf{8} & 7 & \mathbf{7} & 7 \\ & 8 & \mathbf{8} & 8 & \mathbf{7} & 7 & \mathbf{7} \\ & & 7 & \mathbf{7} & 7 & \mathbf{7} & 7 \\ & & & 6 & \mathbf{6} & 6 & \mathbf{5} \\ & & & & 6 & \mathbf{5} & 4 \\ & & & & & 4 & \mathbf{3} \\ & & & & & & 1 \end{array}$$

By Theorem 4.2, we can reduce the properties of ρ to those of the corresponding involution $\tilde{\rho} : \mathcal{P}_{n,m} \rightarrow \mathcal{P}_{n,m}$ defined by

$$\tilde{\rho} = \tilde{\pi}_2\tilde{\pi}_4\cdots \tag{4.2}$$

where the product is over all $\tilde{\pi}_i$ with i even and $\leq n$. In other words, given a plane partition c in $\mathcal{P}_{n,m}$, $\tilde{\rho}$ swaps 1's and 2's in c by the TBK involution, then swap 3's and 4's in c and so on. The resulting plane partition does not depend on the order of the swaps, and is an element of $\mathcal{P}_{n,m}$. Let $\mathcal{P}_{n,m}^{\tilde{\rho}}$ denote the set of elements of $\mathcal{P}_{n,m}$ which is invariant under $\tilde{\rho}$, i.e. $\mathcal{P}_{n,m}^{\tilde{\rho}} = \{c \in \mathcal{P}_{n,m} \mid \tilde{\rho}(c) = c\}$. For example,

if $n = 8$, then the following RCSPP in \mathcal{P}_8 is invariant under $\tilde{\rho}$:

7	4	4	3	2	1	1
6	3	2	1			
5	2					
2	1					
1						

Thus $\mathcal{P}_1^{\tilde{\rho}} = \{\emptyset\}$, $\mathcal{P}_2^{\tilde{\rho}} = \{\emptyset, \boxed{1}\}$, and $\mathcal{P}_3^{\tilde{\rho}}$ is composed of the following 3 RCSPPs:

\emptyset	$\boxed{2}$ $\boxed{1}$	$\boxed{2}$ $\boxed{1}$ $\boxed{1}$
-------------	----------------------------	--

$\mathcal{P}_4^{\tilde{\rho}}$ is composed of the following 10 elements:

\emptyset	$\boxed{2}$ $\boxed{1}$	$\boxed{2}$ $\boxed{1}$ $\boxed{1}$	$\boxed{2}$ $\boxed{1}$	$\boxed{2}$ $\boxed{2}$ $\boxed{1}$ $\boxed{1}$
$\boxed{2}$ $\boxed{2}$ $\boxed{1}$ $\boxed{1}$ $\boxed{1}$	$\boxed{3}$	$\boxed{3}$ $\boxed{2}$ $\boxed{1}$	$\boxed{3}$ $\boxed{2}$ $\boxed{2}$ $\boxed{1}$ $\boxed{1}$	$\boxed{3}$ $\boxed{2}$ $\boxed{1}$ $\boxed{2}$ $\boxed{1}$ $\boxed{1}$

$\mathcal{P}_5^{\tilde{\rho}}$ has 25 elements, and $\mathcal{P}_6^{\tilde{\rho}}$ has 140 elements.

Also in [17, pp.286] the involution $\gamma = \pi_1\pi_3\pi_5 \cdots$ on \mathcal{B}_n is defined and conjectured to have the same effect as the flip of an alternating matrix around the vertical axis (Conjecture 1.2). Naturally we can generalize this definition to the involution $\gamma : \mathcal{B}_{n,m} \rightarrow \mathcal{B}_{n,m}$, $m = 0, 1$, defined by

$$\gamma = \pi_1\pi_3\pi_5 \cdots \tag{4.3}$$

where the product is over all π_i with i odd and $\leq n$. Let $\mathcal{B}_{n,m}^\gamma$ denote the set of elements of $\mathcal{B}_{n,m}$ invariant under γ . For instance, the following TSPP in \mathcal{B}_7 is invariant under γ :

7	7	7	7	7	7
	6	5	5	5	5
		5	5	5	5
			5	5	4
				4	3
					2

The corresponding involution $\tilde{\gamma} : \mathcal{P}_{n,m} \rightarrow \mathcal{P}_{n,m}$, $m = 0, 1$, is defined by

$$\tilde{\gamma} = \tilde{\pi}_1\tilde{\pi}_3\tilde{\pi}_5 \cdots \tag{4.4}$$

where the product is over all $\tilde{\pi}_i$ with i odd and $\leq n$. For $m = 0, 1$, let $\mathcal{P}_{n,m}^{\tilde{\gamma}}$ denote the set of elements of $\mathcal{P}_{n,m}$ invariant under $\tilde{\gamma}$, i.e. $\mathcal{P}_{n,m}^{\tilde{\gamma}} = \{c \in \mathcal{P}_{n,m} \mid \tilde{\gamma}(c) = c\}$. But, since $\mathcal{P}_{n,1} = \mathcal{P}_{n+1,0}$ which implies $\mathcal{P}_{n,1}^{\tilde{\gamma}} = \mathcal{P}_{n+1,0}^{\tilde{\gamma}}$, we only need to study $\mathcal{P}_n^{\tilde{\gamma}}$. Also note that $\mathcal{P}_n^{\tilde{\gamma}} = \emptyset$ unless n is odd. In fact, for $c \in \mathcal{P}_{n,m}$, there are, in total,

exactly $n - 1$ positions where one can put 1's. But, if c is invariant under $\tilde{\gamma}$, then those positions must be half filled. Further one can easily see that the shape of $c_{\geq 2}$ must be even. For example, if $n = 7$, the following RCSPP in \mathcal{P}_7 is invariant under $\tilde{\gamma}$.

5	5	3	2	1
4	4	1		
3	3			
2	2			
1				

Thus we have $\mathcal{P}_3^{\tilde{\gamma}} = \{ \boxed{1} \}$, $\mathcal{P}_5^{\tilde{\gamma}}$ is composed of the following 3 RCSPPs:

1	1		3	2	1		3	3	1
			1				2	2	
							1		

and $\mathcal{P}_5^{\tilde{\gamma}}$ has 26 elements.

By Theorem 4.2, we have established a bijection between $\mathcal{B}_{n,m}^\rho$ and $\mathcal{P}_{n,m}^{\tilde{\rho}}$ and a bijection between $\mathcal{B}_{n,m}^\gamma$ and $\mathcal{P}_{n,m}^{\tilde{\gamma}}$. By this bijection the weight function \bar{U}_r on $\mathcal{B}_{n,m}$ is exactly the same as \bar{U}_r on $\mathcal{P}_{n,m}$. Thus we study $\mathcal{P}_{n,m}^{\tilde{\rho}}$ and $\mathcal{P}_{n,m}^{\tilde{\gamma}}$ with this weight \bar{U}_r in the rest of this paper.

Proof of Theorem 4.2. Let $c = (c_{ij})_{1 \leq i \leq n+m-1, 1 \leq j \leq n}$ be a plane partition in $\mathcal{P}_{n,m}$. We set $b = (b_{ij})_{1 \leq i \leq j \leq n+m-1} = \varphi_{n,m}(c) \in \mathcal{B}_{n,m}$ to be the TSPP mapped by the bijection and set $b' = (b'_{ij})_{1 \leq i \leq j \leq n+m-1} = \pi_r(b) \in \mathcal{B}_{n,m}$ to be the flipped result of b . Let $c' = (c'_{ij})_{1 \leq i \leq n+m-1, 1 \leq j \leq n} = \varphi_{n,m}^{-1}(b') \in \mathcal{P}_{n,m}$ be the corresponding RCSPP.

First, assume $2 \leq r \leq n + m$. Then b' is the resulting TSPP of flipping the part $b_{i,i+r-1}$, $i = 1, \dots, n + m - r$, i.e. replacing $b_{i,i+r-1}$ by $b'_{i,i+r-1}$ for $i = 1, \dots, n + m - r$ where

$$b'_{i,i+r-1} + b_{i,i+r-1} = \min(b_{i-1,i+r-1}, b_{i,i+r-2}) + \max(b_{i,i+r}, b_{i+1,i+r-1}).$$

If we restate this operation $\pi_r : b \mapsto b'$ by the bijection rule (3.4), then this corresponds to the following operation $\tilde{\pi}_r : c \mapsto c'$:

(i) $\theta_i(c'_{\geq r})$ is given by

$$\theta_1(c'_{\geq r}) + \theta_1(c_{\geq r}) = \max(\theta_1(c_{\geq r+1}), \theta_2(c_{\geq r-1})) + \min(n + m - r, \theta_1(c_{\geq r-1}))$$

if $i = 1$, and

$$\theta_i(c'_{\geq r}) + \theta_i(c_{\geq r}) = \max(\theta_i(c_{\geq r+1}), \theta_{i+1}(c_{\geq r-1})) + \min(\theta_{i-1}(c_{\geq r+1}), \theta_i(c_{\geq r-1}))$$

if $i = 2, \dots, n + m - r$.

(ii) If $k \neq r$, then $\theta_i(c'_{\geq k}) = \theta_i(c_{\geq k})$ for all i .

This means the operation $\tilde{\pi}_r$ changes the number of the letters r 's and $r-1$'s in each row i of c , and keeps other letters invariant. If one inspects these rules carefully, then these rules gives precisely the TBK involution.

Next, assume $r = 1$ and $m = 0, 1$. Then b' is the resulting TSPP of replacing b_{ii} by b'_{ii} for $i = 1, \dots, n + m - 1$ where

$$b'_{ii} + b_{ii} = b_{i-1,i} + b_{i,i+1}.$$

Using the rule (3.4) again, the operation $\pi_r : b \mapsto b'$ corresponds to the operation $\tilde{\pi}_r : c \mapsto c'$ with $\theta_i(c'_{\geq r})$ given by

$$\theta_1(c'_{\geq 1}) + \theta_1(c_{\geq 1}) = \theta_1(c_{\geq 2}) + n + m - 1$$

if $i = 1$, and

$$\theta_i(c'_{\geq 1}) + \theta_i(c_{\geq 1}) = \theta_i(c_{\geq 2}) + \theta_{i-1}(c_{\geq 2})$$


if $i = 2, \dots, n + m - 1$. This is exactly the operation $\tilde{\pi}_1 : c \mapsto c'$ we defined above. This completes the proof. \square

5 Twisted domino plane partitions

In this section we consider the invariants of the involution $\tilde{\rho}$ defined in the previous section, which leads us to define a notion of generalized domino plane partitions. In fact, here, we consider a natural bijection as follows. Assume $c \in \mathcal{P}_{n,m}$ is invariant under $\tilde{\rho}$, i.e. invariant under $\tilde{\pi}_{2i}$ for any i . Then we replace paired $2i$ and $2i-1$ in a column by a domino, and replace non-paired k $2i$'s and k $2i-1$'s in a row by k dominoes. But there may remain some saturated parts unchanged, thus we obtain a “twisted” domino plane partition by this operation. In this manner we give a natural bijection of $\mathcal{P}_{n,m}^{\tilde{\rho}}$ to a new object which we denote by $\mathcal{G}_{n,m}$. But we don't know how to count this set at this point. Anyway let's start with definitions.

A *domino* is a special kind of skew shape consists of two squares. A 1×2 domino is called a *horizontal domino* while a 2×1 domino is called a *vertical domino*. Let λ be a partition. A *generalized domino plane partition of shape λ* consists of a tiling of the shape λ by means of ordinary 1×1 squares and dominoes, and a filling of each square or domino with a positive integer so that the integers are weakly decreasing along either rows or columns. The integers in the squares or dominoes are called *parts*. In this paper we call a part a *single part* if it is in a square, i.e. not in a domino. Further we call it a *domino plane partition* if the shape λ is tiled with only dominoes, i.e. without a single square. We say that a part is in the i th row (resp. j th column) if the square or domino which contains the number intersects with the i th row (resp. j th column) of λ . A (generalized) domino plane partition is said to be *column-strict* if it is strictly decreasing along each column. For example, the left-below is a column-strict generalized domino plane partition of shape $(4, 3, 2, 1)$,

while the right-below is a column-strict domino plane partition of shape $(4, 4, 2)$.




Definition 5.1. Let m and $n \geq 1$ be nonnegative integers. Let $\mathcal{G}_{n,m}$ denote the set of column-strict generalized domino plane partitions c subject to the constraints that


- (E1) c has at most n columns;
- (E2) each part in the j th column does not exceed $\lceil (n + m - j)/2 \rceil$;
- (E3) If a part in the j th column is equal to $\lceil (n + m - j)/2 \rceil$ and $n + m - j$ is odd, then it must be a single part. On the other hand, if a single part appears in the j th column of c , then $n + m - j$ must be odd and it must be equal to $\lceil (n + m - j)/2 \rceil$.

We call an element in $\mathcal{G}_{n,m}$ a *twisted domino plane partition*, and we simply write \mathcal{G}_n for $\mathcal{G}_{n,0}$. If c is in $\mathcal{G}_{n,m}$, we call a single part in c a *saturated* part, which can appear only in the first row. Thus a saturated part is always a single part and vice versa, which equals $\lceil (n + m - j)/2 \rceil$ appearing in the cell $(1, j)$ where $n + m - j$ is odd.

For instance, $\mathcal{G}_1 = \{\emptyset\}$, $\mathcal{G}_2 = \{\emptyset, \boxed{1}\}$, \mathcal{G}_3 is composed of the following 3 elements:



\mathcal{G}_4 is composed of the following 10 elements:



\mathcal{G}_5 has 25 elements and \mathcal{G}_6 has 140 elements. For example, if $c \in \mathcal{G}_5$, all parts in the 1st and 2nd columns are ≤ 2 , and all parts in the 3rd and 4th columns are ≤ 1 . A saturated (=single) part equal to 2 can appear in the cell $(1, 2)$, and a saturated part equal to 1 can appear in the cell $(1, 4)$.

For $c \in \mathcal{G}_{n,m}$, let $\bar{U}_1(c)$ denote the number of 1's in c . From the above examples we obtain that the first few terms of $\sum_{c \in \mathcal{G}_r} t^{\bar{U}_1(c)}$ are $1, 1+t, 1+t+t^2, (1+t)(2+t+2t^2), 3+6t+7t^2+6t^3+3t^4$ and $5(1+t)(1+t^2)(2+3t+2t^2)$.

Let m and $n \geq 1$ be nonnegative integers. Let's consider the situation when we apply $\tilde{\pi}_r$ to a plane partition c in $\mathcal{P}_{n,m}$. We say an entry r or $r - 1$ in c is *free* if it is not saturated $r - 1$ nor there is no corresponding $r - 1$ or r in the same column. If r and $r - 1$ are in the same column, we say they are *paired*. Thus an $r - 1$ in c can be free, paired or saturated, whereas an r in c can be free or paired.

Let c be a plane partition in $\mathcal{P}_{n,m}^{\tilde{\rho}}$, i.e. c is invariant under each $\tilde{\pi}_{2r}$ for $r \geq 1$. We associate to c a column-strict generalized domino plane partition d as follows. For each $r \geq 1$, we replace paired $2r$ and $2r - 1$ by a vertical domino containing r , and, if row i contains k free $2r$'s and k free $2r - 1$'s, then we replace these with k horizontal dominoes containing r . Finally a saturated part equal to $2r - 1$ should be replaced by a single part r . Let us denote by $\psi_{n,m}(c)$ the resulting generalized domino plane partition d . We may use the abbreviated notation $\psi_n(c)$ for $\psi_{n,0}(c)$. For example, the left-below plane partition in $\mathcal{P}_8^{\tilde{\rho}}$ is mapped to the right-below generalized plane partition in \mathcal{G}_8 by ψ_8 :

7	4	4	3	2	1	1
6	3	2	1			
5	2					
2	1					
1						

4	2	2	1	1
3		1		
	1			
1				

Theorem 5.2. Let m and $n \geq 1$ be nonnegative integers and let $c = (c_{ij})_{1 \leq i \leq n+m, 1 \leq j \leq n}$ be a plane partition in $\mathcal{P}_{n,m}^{\tilde{\rho}}$. Associate to c the generalized domino plane partition $\psi_{n,m}(c)$ as above. Then $\psi_{n,m}(c)$ is in $\mathcal{G}_{n,m}$, and $\psi_{n,m}$ is a bijection between $\mathcal{P}_{n,m}^{\tilde{\rho}}$ and $\mathcal{G}_{n,m}$.

6 Domino plane partitions

In this section, we introduce two important classes of domino plane partitions, i.e. $\mathcal{D}_{n,m}^R$ and $\mathcal{D}_{n,m}^C$. The main result of this section is Theorem 6.2 which shows that there is a bijection between the set $\mathcal{P}_{2n+1}^{\tilde{\gamma}}$ of restricted column-strict plane partitions invariant under $\tilde{\gamma}$ and the set \mathcal{D}_{2n-1}^R of restricted domino plane partitions with all rows of even length. At the end of this section we state a conjecture that the set $\mathcal{G}_{n,m}$ of twisted domino plane partitions and the set $\mathcal{D}_{n,m}^C$ of restricted domino plane partitions with all columns of even length would have the same cardinality, and the statistics \bar{U}_1 would have the same distribution. Now we start from the definition of these classes.

Definition 6.1. Let m and $n \geq 1$ be nonnegative integers. Let $\mathcal{D}_{n,m}$ denote the set of column-strict domino plane partitions $d = (d_{ij})_{1 \leq i,j}$ subject to the constraints that

(D1) d has at most n columns;

(D2) each part in the j th column does not exceed $\lceil (n + m - j)/2 \rceil$;

An element of $\mathcal{D}_{n,m}$ is called a *restricted domino plane partition* (abbreviated to RDPP). If a part in the j th column of c is equal to $\lceil (n+m-j)/2 \rceil$, we call the part *saturated*. For $d \in \mathcal{D}_{n,m}$ and a positive integer $r \geq 1$, let $\bar{U}_r(d)$ denote the number of parts equal to r plus the number of saturated parts less than r . Further, if d in $\mathcal{D}_{n,m}$ satisfy the condition that

(D3) each row (resp. column) of d has even length,

then d is called a restricted column-strict domino plane partition *with all rows (resp. columns) of even length*. The set of all $d \in \mathcal{D}_{n,m}$ with all rows (resp. columns) of even length is denoted by $\mathcal{D}_{n,m}^R$ (resp. $\mathcal{D}_{n,m}^C$). When $m = 0$, we write \mathcal{D}_n for $\mathcal{D}_{n,0}$, \mathcal{D}_n^R for $\mathcal{D}_{n,0}^R$ and \mathcal{D}_n^C for $\mathcal{D}_{n,0}^C$.

For example, $\mathcal{D}_1^R = \mathcal{D}_2^R = \{\emptyset\}$, \mathcal{D}_3^R is composed of the following 3 elements:

$$\emptyset, \quad \boxed{1}, \quad \begin{array}{|c|c|} \hline 1 & 1 \\ \hline \end{array}.$$

\mathcal{D}_4^R is composed of the following 4 elements:

$$\emptyset, \quad \boxed{1}, \quad \begin{array}{|c|c|} \hline 1 & 1 \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline 2 & 1 \\ \hline \end{array}.$$

\mathcal{D}_5^R has 26 elements, \mathcal{D}_6^R has 50 elements, and \mathcal{D}_7^R has 646 elements.

Let n be a positive integer, and assume $c = (c_{ij})_{1 \leq i, j \leq n}$ is in $\mathcal{P}_{2n+1}^{\tilde{\gamma}}$, i.e. c is invariant under each $\tilde{\pi}_{2r-1}$ for $r \geq 1$. For each $\tilde{\pi}_{2r-1}$, we use the notation “free”, “paired” and “saturated” for parts equal to $2r+1$ or $2r$ as before. We associate to c a column-strict (generalized) domino plane partition d as follows. First of all, since c is invariant under $\tilde{\pi}_1$, which means the shape of $c_{\geq 2}$ must be even. Thus we remove all 1’s from c so that each row of the resulting plane partition $c_{\geq 2}$ has even length. Next, since $c_{\geq 2}$ is invariant under $\tilde{\pi}_3$, we replace paired 3 and 2 by a vertical domino containing 1, and, if row i contains k free 3’s followed by k free 2’s, then we replace them by k horizontal dominoes containing 1. If there exists a saturated part equal to 2, it should be replaced by a single box containing 1. Next we replace paired 5 and 4 by a vertical domino containing 2, and, if row i contains k free 5’s followed by k free 4’s, then we replace them with k horizontal dominoes containing 2. If there is a saturated part equal to 4, it is replaced by a single box containing 2. We repeat this process and finally obtain a column-strict (generalized) domino plane partition d with all rows of even length. Let us denote the resulting (generalized) domino plane partition d by $\tau_{2n+1}(c)$. For example, the left-below plane partition c in $\mathcal{P}_{11}^{\tilde{\gamma}}$ is mapped to the right-below generalized domino plane partition d in \mathcal{G}_9 by τ_{11} :

$$c = \begin{array}{|c|c|c|c|c|c|c|c|} \hline 7 & 7 & 6 & 6 & 3 & 2 & 1 & 1 \\ \hline 5 & 5 & 4 & 3 & 1 & & & \\ \hline 4 & 3 & 2 & 2 & & & & \\ \hline 1 & 1 & & & & & & \\ \hline \end{array} \quad d = \begin{array}{|c|c|c|} \hline 3 & 3 & 1 \\ \hline 2 & 2 & 1 \\ \hline & 1 & \\ \hline \end{array}$$

Theorem 6.2. Let n be a positive integer. Let c be a plane partition in $\mathcal{P}_{2n+1}^{\tilde{\gamma}}$. Then, $\tau_{2n+1}(c)$ has no saturated part, i.e. $\tau_{2n+1}(c) \in \mathcal{D}_{2n-1}^{\text{R}}$. Thus τ_{2n+1} gives a bijection of $\mathcal{P}_{2n+1}^{\tilde{\gamma}}$ onto $\mathcal{D}_{2n-1}^{\text{R}}$. Further we have $\overline{U}_1(\tau_{2n+1}(c)) = \overline{U}_2(c)$.

Proof. Let c be a plane partition in $\mathcal{P}_{2n+1}^{\tilde{\gamma}}$ and let $d = \tau_{2n+1}(c)$. Note that a saturated part $n+1-i$ ($i = 1, \dots, n$) in d can appear in the cell $(1, 2i-1)$ if there exists. Let λ be the shape of $c_{\geq 2}$. As a tableau can be expressed by a sequence of partitions (see [15, p.5]), we can write the generalized domino plane partition d by a sequence

$$\emptyset = \lambda^{(0)} \subseteq \lambda^{(1)} \subseteq \dots \subseteq \lambda^{(n)} = \lambda$$

of partitions in which $\lambda^{(i)}/\lambda^{(i-1)}$ consists of all the cells and dominoes that contain the letter $n+1-i$. If necessary, we add a zero at the end of λ and we write $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_{2r})$ as a sequence of even length. Then we can write $\lambda^{(i)} = (\lambda_1^{(i)}, \dots, \lambda_{2r}^{(i)})$ with $\lambda_1^{(i)} \geq \dots \geq \lambda_{2r}^{(i)} \geq 0$ for $i = 1, \dots, n$. For example, the above domino tableau d in \mathcal{G}_9 is expressed by the sequence: $\lambda^{(0)} = (0, 0, 0, 0)$, $\lambda^{(1)} = (4, 0, 0, 0)$, $\lambda^{(2)} = (4, 3, 1, 0)$ and $\lambda^{(3)} = (6, 4, 4, 0)$. If we put $\ell_j^{(i)} = \lambda_j^{(i)} + 2r - j$ for $i = 1, \dots, n$ and $j = 1, \dots, 2r$, then we have $\ell_1^{(i)} > \dots > \ell_{2r}^{(i)} \geq 0$ for $i = 1, \dots, n$. In the above example, we have $\ell^{(0)} = (3, 2, 1, 0)$, $\ell^{(1)} = (7, 2, 1, 0)$, $\ell^{(2)} = (7, 5, 2, 0)$ and $\ell^{(3)} = (9, 6, 5, 0)$. When $i = 0$, we have $\ell_j^{(0)} = 2r - j$ for $j = 1, \dots, 2n$, and $\ell^{(0)}$ consist of r even integers and r odd integers. For each $i = 1, \dots, n$, if there is no saturated part equal to $n+1-i$, then $\lambda^{(i)}/\lambda^{(i-1)}$ contains only dominoes, thus the cardinalities of odd integers and even integers in $\ell^{(i)}$ are the same as those of $\ell^{(i-1)}$. Assume there was certain i such that $\lambda^{(i)}/\lambda^{(i-1)}$ contain a saturated part $n+1-i$ in $(1, 2i-1)$. Then the cardinality of odd integer would decrease by 1, and the cardinality of even integer would increase by 1 if we compare $\ell^{(i)}$ with $\ell^{(i-1)}$. Thus, if there were saturated parts, finally $\ell^{(n)}$ would contain less odd integers than even integers. Since λ is even partition, we have $\ell_j^{(n)} + j$ must be even for $j = 1, \dots, 2r$, which implies we must have the same number of even integers and odd integers in $\ell^{(n)}$. This is a contradiction. Thus we conclude that d has no saturated part and it is easy to see that d is in $\mathcal{D}_{2n-1}^{\text{R}}$. The construction of $d \in \mathcal{D}_{2n-1}^{\text{R}}$ from $c \in \mathcal{P}_{2n+1}^{\tilde{\gamma}}$ is clearly reversible and give a bijection of $\mathcal{P}_{2n+1}^{\tilde{\gamma}}$ onto $\mathcal{D}_{2n-1}^{\text{R}}$. Since 3's and 2's in c is replaced by dominoes containing 1's, we have $\overline{U}_2(c) = \overline{U}_1(d)$. This completes the proof. \square

Next we give examples of \mathcal{D}_k^{C} . We have $\mathcal{D}_1^{\text{C}} = \{\emptyset\}$, $\mathcal{D}_2^{\text{C}} = \left\{ \emptyset, \begin{array}{|c|} \hline \mathbf{1} \\ \hline \end{array} \right\}$, and \mathcal{D}_3^{C} has the following 3 elements:

$$\emptyset, \quad \begin{array}{|c|} \hline \mathbf{1} \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline \mathbf{1} & \mathbf{1} \\ \hline \end{array}.$$

\mathcal{D}_4^{C} has the following 10 elements:

$$\emptyset, \quad \begin{array}{|c|} \hline \mathbf{1} \\ \hline \end{array}, \quad \begin{array}{|c|} \hline \mathbf{2} \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline \mathbf{1} & \mathbf{1} \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline \mathbf{2} & \mathbf{1} \\ \hline \end{array},$$

$$\begin{array}{|c|c|c|} \hline 1 & \mathbf{1} & \mathbf{1} \\ \hline \end{array}, \quad
\begin{array}{|c|c|c|} \hline \mathbf{2} & \mathbf{1} & \mathbf{1} \\ \hline \end{array}, \quad
\begin{array}{|c|} \hline \mathbf{2} \\ \hline 1 \\ \hline \end{array}, \quad
\begin{array}{|c|c|} \hline \mathbf{2} & \mathbf{1} \\ \hline 1 & \\ \hline \end{array}, \quad
\begin{array}{|c|c|c|} \hline \mathbf{2} & \mathbf{1} & \mathbf{1} \\ \hline 1 & & \\ \hline \end{array},$$

and it is not hard to see that \mathcal{D}_5^C has 25 elements, \mathcal{D}_6^C has 140 elements, and \mathcal{D}_7^C has 588 elements. In this example the bold-faced parts are saturated. The author computed examples for small n , m and observe that the cardinalities of $\mathcal{G}_{n,m}$ and $\mathcal{D}_{n,m}^C$ agree for $n \leq 6$.

Conjecture 6.3. Let m and $n \geq 1$ be non-negative integers. Then there would be a bijection which proves that $\mathcal{G}_{n,m}$ and $\mathcal{D}_{n,m}^C$ has the same cardinality. Moreover this bijection keeps \bar{U}_1 (i.e. the number of 1s) invariant.

7 Determinantal formulae

A standard map which associate a k -tuple of tableaux with a given k -rim hook tableaux is presented in the paper [25, Section 6] (see also [7, Theorem 6.3]). Essentially we can use this map to associate a pair of column-strict plane partitions in $\mathcal{P}_{n,m}$ with a domino plane partition in $\mathcal{D}_{n,m}$. By this map we can rewrite the statistics \bar{U}_k on $\mathcal{D}_{n,m}$ defined in the previous section as the sum of the statistics of each column-strict plane partition. Thus we can obtain the generating functions of $\mathcal{D}_{n,m}^R$ and $\mathcal{D}_{n,m}^C$ as an application of [8, Lemma 7.1] (see Theorem 7.2, Corollary 7.7). As a corollary of these generating functions we obtain a determinantal formula (Corollary 7.8(ii)) for Conjecture 1.2. Thus Conjecture 1.2 reduce to a determinant evaluation problem (Conjecture 7.9(i)). In the special case where $t = 1$, we prove the conjecture from Andrews-Burge determinant (see Lemma 7.10, Theorem 7.11).

Let λ be a partition. We define a pair $(\lambda^{(0)}, \lambda^{(1)})$ which is called *2-quotient* of λ as follows. If necessary, we add a zero at the end of λ and we regard $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_{2n})$ as a sequence of integers of even length. Let $\ell = \lambda + \delta_{2n} = (\lambda_1 + 2n - 1, \lambda_2 + 2n - 2, \dots, \lambda_{2n})$, and put $\ell = \ell^{(0)} \uplus \ell^{(1)}$ where $\ell^{(0)} = \{x \in \ell : x \equiv 0 \pmod{2}\}$ and $\ell^{(1)} = \{x \in \ell : x \equiv 1 \pmod{2}\}$. We can write $\ell^{(0)} = (2k_1^{(0)}, \dots, 2k_r^{(0)})$ and $\ell^{(1)} = (2k_1^{(1)} + 1, \dots, 2k_s^{(1)} + 1)$ where $k_1^{(0)} > \dots > k_r^{(0)} \geq 0$ and $k_1^{(1)} > \dots > k_s^{(1)} \geq 0$. The partition $\lambda^{(0)}$ (resp. $\lambda^{(1)}$) is defined to be $(k_1^{(0)} - r + 1, k_2^{(0)} - r + 2, \dots, k_r^{(0)})$ (resp. $(k_1^{(1)} - s + 1, k_2^{(1)} - s + 2, \dots, k_s^{(1)})$). For example, if $\lambda = (5, 5, 3, 1, 1, 1)$, then we have $\lambda^{(0)} = (3, 2, 1)$ and $\lambda^{(1)} = (2)$.

Definition 7.1. Let m and $n \geq 1$ be nonnegative integers. Let $\mathcal{Q}_{n,m}$ denote the set of all pairs $p = (c_0, c_1)$ of plane partitions such that

- (i) $c_0 \in \mathcal{P}_{n_0, m_0}$ where $n_0 = \lfloor \frac{n}{2} \rfloor$ and $m_0 = \lfloor \frac{n+m+1}{2} \rfloor - n_0$,
- (ii) $c_1 \in \mathcal{P}_{n_1, m_1}$ where $n_1 = \lfloor \frac{n}{2} \rfloor$ and $m_1 = \lfloor \frac{n+m+1}{2} \rfloor - n_1$.

For $p \in \mathcal{Q}_{n,m}$ and a positive integer $r \geq 1$, let $\bar{U}_r(p) = \bar{U}_r(c_0) + \bar{U}_r(c_1)$. Further let $\mathcal{Q}_{n,m}^V$ denote the set of all pairs $p = (c_0, c_1) \in \mathcal{Q}_{n,m}$ such that $\text{sh}(c_1) \subseteq \text{sh}(c_0)$ and $\text{sh}(c_0) \setminus \text{sh}(c_1)$ is a vertical strip. Meanwhile, let $\mathcal{Q}_{n,m}^H$ denote the set of all pairs

$p = (c_0, c_1) \in \mathcal{Q}_{n,m}$ such that $\text{sh}(c_0) \subseteq \text{sh}(c_1)$ and $\text{sh}(c_1) \setminus \text{sh}(c_0)$ is a horizontal strip. We also write \mathcal{Q}_n^V for $\mathcal{Q}_{n,0}^V$, and \mathcal{Q}_n^H for $\mathcal{Q}_{n,0}^H$ in short.


For example, if $n = 4$ and $m = 0$, then the pairs (c_0, c_1) in \mathcal{Q}_4^V satisfy the condition that $c_0 \in \mathcal{P}_{2,1}$, $c_1 \in \mathcal{P}_2$ and $\text{sh}(c_0) \setminus \text{sh}(c_1)$ is a vertical strip. Thus, \mathcal{Q}_4^V has the following 10 elements:

$$(\emptyset, \emptyset), \quad (\boxed{1}, \emptyset), \quad (\boxed{1}, \boxed{1}), \quad (\boxed{11}, \boxed{1}), \quad (\boxed{2}, \emptyset), \\ (\boxed{2}, \boxed{1}), \quad (\boxed{21}, \boxed{1}), \quad (\boxed{2}, \emptyset), \quad (\boxed{2}, \boxed{1}), \quad (\boxed{21}, \boxed{1}).$$


The italic characters stand for saturated parts. Meanwhile, the pairs (c_0, c_1) in \mathcal{Q}_4^H satisfy the condition that $c_0 \in \mathcal{P}_{2,1}$, $c_1 \in \mathcal{P}_2$ and $\text{sh}(c_1) \setminus \text{sh}(c_0)$ is a horizontal strip. Thus, \mathcal{Q}_4^H has the following 4 elements:

$$(\emptyset, \emptyset), \quad (\emptyset, \boxed{1}), \quad (\boxed{1}, \boxed{1}), \quad (\boxed{2}, \boxed{1}).$$

Here we describe a bijection of column-strict domino plane partitions onto pairs of column-strict plane partitions. Given a domino α , the upper-rightmost cell in α is called the *head* of α , and the lower-leftmost cell in α is called the *tail* of α . For any skew-shape, we number the diagonals, beginning with 0 for the main diagonal, increasing up and to the right and decreasing down and to the left. The *diagonal diag* (α) of a domino α is the diagonal of the head of α , and the *color* $\text{Color}(\alpha)$ of α is $\text{diag}(\alpha) \bmod 2$. In the following pictures, we draw only the diagonals whose numbers are even.



Assume we are given a column-strict domino plane partition d . For each $k = 0, 1$, we associate a column-strict plane partition c_k with d . Along each diagonal $\text{diag}(r)$, we read only the numbers in the dominoes that cross $\text{diag}(r)$ and has color k . We replace the dominoes by single cells containing the numbers and slide down along the diagonal $\text{diag}(r)$. In this way we obtain an (ordinary) column-strict partition c_k . For example, we associate the column-strict domino plane partition



the pair

$$c_0 = \begin{array}{|c|c|} \hline 1 & 1 \\ \hline \end{array} \quad c_1 = \begin{array}{|c|c|c|} \hline 3 & 3 & 1 \\ \hline 2 & 2 & \\ \hline \end{array}$$

of plane partitions. Let Φ denote the map which associate the pair (c_0, c_1) of column-strict plane partitions with a column-strict domino plane partition d (cf. [25, Section 6], [7, Theorem 6.3]).

Theorem 7.2. The restriction of the map $\Phi : d \mapsto (c_0, c_1)$ to $\mathcal{D}_{n,m}$ gives a bijection of $\mathcal{D}_{n,m}$ onto $\mathcal{Q}_{n,m}$. By this bijection the $(\text{sh}(c_0), \text{sh}(c_1))$ is the 2-quotient of $\text{sh}(d)$, and we have $\overline{U}_r(d) = \overline{U}_r(\Phi(d))$ for $d \in \mathcal{D}_{n,m}$ and $r \geq 1$. Especially, the restriction of the map Φ to $\mathcal{D}_{n,m}^{\text{R}}$ (resp. $\mathcal{D}_{n,m}^{\text{C}}$) gives a bijection of $\mathcal{D}_{n,m}^{\text{R}}$ (resp. $\mathcal{D}_{n,m}^{\text{C}}$) onto $\mathcal{Q}_{n,m}^{\text{H}}$ (resp. $\mathcal{Q}_{n,m}^{\text{V}}$) which preserve the statistics \overline{U}_r .

Proof. The first half of the theorem is an easy consequence of the definition. The latter half of the theorem follows from the following proposition. \square

Proposition 7.3. Let d be a column-strict domino plane partition, and let $(c_0, c_1) = \Phi(d)$. Then

- (i) All rows of d have even length if, and only if, $\text{sh}(c_0) \subseteq \text{sh}(c_1)$ and $\text{sh}(c_1) \setminus \text{sh}(c_0)$ is a horizontal strip.
- (ii) All columns of d have even length if, and only if, $\text{sh}(c_1) \subseteq \text{sh}(c_0)$ and $\text{sh}(c_0) \setminus \text{sh}(c_1)$ is a vertical strip.
- (iii) All rows and columns of d have even length if, and only if, $\text{sh}(c_0) = \text{sh}(c_1)$.

Proof. Let $\lambda = \text{sh}(d)$ and let $(\lambda^{(0)}, \lambda^{(1)})$ denote the 2-quotient of λ . We regard $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_{2n})$ as a sequence of integers of even length as before. Put $\ell = \lambda + \delta_{2n}$, and let $\ell^{(0)} = \{x \in \ell : x \equiv 0 \pmod{2}\}$ and $\ell^{(1)} = \{x \in \ell : x \equiv 1 \pmod{2}\}$.

- (i) Note that λ is even partition if, and only if, $\ell_i \equiv i \pmod{2}$. This implies $\ell^{(0)} = (\ell_2, \dots, \ell_{2n}) = (2k_1^{(0)}, \dots, 2k_n^{(0)})$ and $\ell^{(1)} = (\ell_1, \dots, \ell_{2n-1}) = (2k_1^{(1)} + 1, \dots, 2k_n^{(1)} + 1)$ with $k_1^{(1)} \geq k_1^{(0)} > k_2^{(1)} \geq k_2^{(0)} > \dots > k_n^{(1)} \geq k_n^{(0)} \geq 0$. Thus, if we put $\lambda^{(0)} = (\lambda_1^{(0)}, \dots, \lambda_n^{(0)})$ and $\lambda^{(1)} = (\lambda_1^{(1)}, \dots, \lambda_n^{(1)})$, then we have $\lambda_1^{(1)} \geq \lambda_1^{(0)} \geq \lambda_2^{(1)} \geq \lambda_2^{(0)} \geq \dots \geq \lambda_n^{(1)} \geq \lambda_n^{(0)} \geq 0$. This proves that $\lambda^{(0)} \subseteq \lambda^{(1)}$ and $\lambda^{(1)} \setminus \lambda^{(0)}$ is a horizontal strip. The reverse can be proved similarly.
- (ii) Note that λ' is even if, and only if, $\ell_{2i-1} = \ell_{2i} + 1$ for $i = 1, \dots, n$. This implies $\ell^{(0)} = (2k_1^{(0)}, \dots, 2k_n^{(0)})$ and $\ell^{(1)} = (2k_1^{(1)} + 1, \dots, 2k_n^{(1)} + 1)$ with $k_1^{(1)} > \dots > k_n^{(1)} \geq 0$, $k_1^{(1)} > \dots > k_n^{(1)} \geq 0$ and $k_i^{(1)} = k_i^{(0)}$ or $k_i^{(0)} - 1$ for $i = 1, \dots, n$. Thus, if we put $\lambda^{(0)} = (\lambda_1^{(0)}, \dots, \lambda_n^{(0)})$ and $\lambda^{(1)} = (\lambda_1^{(1)}, \dots, \lambda_n^{(1)})$, we have $\lambda_i^{(1)} = \lambda_i^{(0)}$ or $\lambda_i^{(0)} - 1$ which is equivalent that $\lambda^{(1)} \subseteq \lambda^{(0)}$ and $\lambda^{(0)} \setminus \lambda^{(1)}$ is a vertical strip.
- (iii) This immediately follows from (i) and (ii).

This completes the proof. \square

Lemma 7.4. Let n and N be positive integers such that $n \leq N$. Let $A = (a_{ij})$ and $B = (b_{ij})$ be $n \times N$ matrices.

- (i) Let $C = (c_{ij})$ denote the $n \times N$ matrix whose (i, j) th entry is $c_{ij} = \sum_{k=1}^j b_{ik}$. Then we have

$$\sum \det \begin{pmatrix} a_{1,j_1} & \cdots & a_{1,j_n} \\ \vdots & \ddots & \vdots \\ a_{n,j_1} & \cdots & a_{n,j_n} \end{pmatrix} \det \begin{pmatrix} b_{1,k_1} & \cdots & b_{1,k_n} \\ \vdots & \ddots & \vdots \\ b_{n,k_1} & \cdots & b_{n,k_n} \end{pmatrix} = \det A^t C, \quad (7.1)$$

where the sum runs over all (j_1, \dots, j_n) and (k_1, \dots, k_n) such that

$$1 \leq k_1 \leq j_1 < k_2 \leq j_2 < \dots < k_n \leq j_n \leq N.$$

- (ii) Let $C' = (c'_{ij})$ denote the $n \times N$ matrix whose (i, j) th entry is $c'_{ij} = \sum_{k=\max(1, j-1)}^j b_{ik}$. Then we have

$$\sum \det \begin{pmatrix} a_{1,j_1} & \dots & a_{1,j_n} \\ \vdots & \ddots & \vdots \\ a_{n,j_1} & \dots & a_{n,j_n} \end{pmatrix} \det \begin{pmatrix} b_{1,k_1} & \dots & b_{1,k_n} \\ \vdots & \ddots & \vdots \\ b_{n,k_1} & \dots & b_{n,k_n} \end{pmatrix} = \det A {}^t C', \quad (7.2)$$

where the sum runs over all (j_1, \dots, j_n) and (k_1, \dots, k_n) such that

$$1 \leq j_1 < j_2 < \dots < j_n \leq N,$$

and $\max(j_{\nu-1} + 1, j_\nu - 1) \leq k_\nu \leq j_\nu$ for $\mu = 1, 2, \dots, n$. Here we use the convention that $j_0 = 1$.

Proof. The both of the identities (7.1), (7.2) reduce to the following Cauchy-Binet identity: Let $X = (x_{ij})$ and $Y = (y_{ij})$ be $n \times N$ matrices. Then

$$\sum \det \begin{pmatrix} x_{1,j_1} & \dots & x_{1,j_n} \\ \vdots & \ddots & \vdots \\ x_{n,j_1} & \dots & x_{n,j_n} \end{pmatrix} \det \begin{pmatrix} y_{1,j_1} & \dots & y_{1,j_n} \\ \vdots & \ddots & \vdots \\ y_{n,j_1} & \dots & y_{n,j_n} \end{pmatrix} = \det X {}^t Y, \quad (7.3)$$

where the sum runs over all $1 \leq j_1 < j_2 < \dots < j_n \leq N$. For example, to prove (7.1) using (7.3), it suffices to show that

$$\det C_{j_1, j_2, \dots, j_n} = \sum_{k_1=1}^{j_1} \sum_{k_2=j_1+1}^{j_2} \dots \sum_{k_n=j_{n-1}+1}^{j_n} B_{k_1, k_2, \dots, k_n}.$$

This can be easily seen if one writes

$$c_{i, j_\nu} = \sum_{k=1}^{j_\nu} b_{i,k} = \sum_{k=1}^{j_1} b_{i,k} + \sum_{k=j_1+1}^{j_2} b_{i,k} + \dots + \sum_{k=j_{\nu-1}+1}^{j_\nu} b_{i,k}$$

for $1 \leq i, \nu \leq n$. The other identity (7.2) can be derived similarly. \square

In [8] we obtained the generating function of $\mathcal{P}_{n,m}$ weighted by the statistics \bar{U}_k and parts. Here we cite the theorem as the following lemma without proof.

Lemma 7.5. ([8, Lemma 7.1]) Let m and $n \geq 1$ be non-negative integers, and fix a positive integer $N \geq n + m$. Let λ be a partition with $\ell(\lambda) \leq n$. For $c \in \mathcal{P}_{n,m}$, let $\mathbf{t}^{\bar{U}(c)} \mathbf{x}^c$ denote $\prod_{k=1}^N t_k^{\bar{U}_k(c)} \prod_{i \geq 1} x_i^{m_i}$, where m_i denote the number of i 's in c . We put $y_i = t_i x_i$ and $Y_i = \prod_{k=i}^N t_k x_i$, and use the vector notation $\mathbf{y}^{(r)}$ for the r -tuples (y_1, \dots, y_r) of the variables. Then the generating function of all plane partitions $c \in \mathcal{P}_{n,m}$ of shape λ' with the weight $\mathbf{t}^{\bar{U}(c)} \mathbf{x}^c$ is given by

$$\sum_{\substack{c \in \mathcal{P}_{n,m} \\ \text{sh}(c) = \lambda'}} \mathbf{t}^{\bar{U}(c)} \mathbf{x}^c = \det \left(e_{\lambda_j - j + i}^{(n+m-i)}(\mathbf{y}^{(n+m-i-1)}, Y_{n+m-i}) \right)_{1 \leq i, j \leq n}. \quad (7.4)$$

As an application of Lemma 7.4 and Lemma 7.5, we obtain the following theorem, which give us the generating function of $\mathcal{D}_{n,m}^R$ and $\mathcal{D}_{n,m}^C$ with the same weights.

Theorem 7.6. Let m and $n \geq 1$ be nonnegative integers, and fix a positive integer $N \geq n + m$. Put $n_0 = \lceil \frac{n}{2} \rceil$, $m_0 = \lceil \frac{n+m+1}{2} \rceil - n_0$, $n_1 = \lfloor \frac{n}{2} \rfloor$ and $m_1 = \lfloor \frac{n+m+1}{2} \rfloor - n_1$. For a domino plane partition $d \in \mathcal{D}_{n,m}$, let $\mathbf{t}^{\bar{U}(d)} \mathbf{x}^d$ denote $\prod_{k=1}^N t_k^{\bar{U}_k(d)} \prod_{i \geq 1} x_i^{m_i}$ where m_i denote the number of i 's in d . We put $y_i = t_i x_i$ and $Y_i = \prod_{k=i}^N t_k x_i$, and write $\mathbf{y}^{(r)}$ for the r -tuples (y_1, \dots, y_r) , where we use the convention that $(\mathbf{y}^{(r-1)}, Y_r)$ is empty if $r \leq 0$.

- (i) If n is even (i.e. $n_0 = n_1$), then we obtain $\sum_{d \in \mathcal{D}_{n,m}^R} \mathbf{t}^{\bar{U}(d)} \mathbf{x}^d = \det R$, where $R = (R_{ij})_{0 \leq i \leq n_0-1, 0 \leq j \leq n_1-1}$ is the $n_0 \times n_1$ matrix whose (i, j) th entry is

$$R_{ij} = \sum_{k \geq 0} e_{k-i}^{(m_0+i+1)}(\mathbf{y}^{(m_0+i-1)}, Y_{m_0+i}, 1) e_{k-j}^{(m_1+j)}(\mathbf{y}^{(m_1+j-1)}, Y_{m_1+j}). \quad (7.5)$$

- (ii) If n is odd (i.e. $n_0 = n_1 + 1$), then we obtain $\sum_{d \in \mathcal{D}_{n,m}^R} \mathbf{t}^{\bar{U}(d)} \mathbf{x}^d = \det(\vec{r} \mid R)$, where

$$R_{ij} = \sum_{k \geq 0} e_{k-i}^{(m_0+i+1)}(\mathbf{y}^{(m_0+i-1)}, Y_{m_0+i}, 1) e_{k-j-1}^{(m_1+j)}(\mathbf{y}^{(m_1+j-1)}, Y_{m_1+j}), \quad (7.6)$$

and $\vec{r} = (r_i)_{0 \leq i \leq n_0-1}$ is the column vector whose i th entry is $r_i = \delta_{i,0}$.

- (iii) If n is even (i.e. $n_0 = n_1$), then we obtain $\sum_{d \in \mathcal{D}_{n,m}^C} \mathbf{t}^{\bar{U}(d)} \mathbf{x}^d = \det C$, where $C = (C_{ij})_{0 \leq i \leq n_0-1, 0 \leq j \leq n_1-1}$ is the $n_0 \times n_1$ matrix whose (i, j) th entry is

$$C_{ij} = \sum_{k \geq 0} \sum_{\nu=0}^k e_{k-i}^{(m_0+i)}(\mathbf{y}^{(m_0+i-1)}, Y_{m_0+i}) e_{\nu-j}^{(m_1+j)}(\mathbf{y}^{(m_1+j-1)}, Y_{m_1+j}). \quad (7.7)$$

- (iv) If n is odd (i.e. $n_0 = n_1 + 1$), then we obtain $\sum_{d \in \mathcal{D}_{n,m}^C} \mathbf{t}^{\bar{U}(d)} \mathbf{x}^d = \det(\vec{c} \mid C)$, where

$$C_{ij} = \sum_{k \geq 0} \sum_{\nu=0}^{k-1} e_{k-i}^{(m_0+i)}(\mathbf{y}^{(m_0+i-1)}, Y_{m_0+i}) e_{\nu-j}^{(m_1+j)}(\mathbf{y}^{(m_1+j-1)}, Y_{m_1+j}), \quad (7.8)$$

and $\vec{c} = (c_i)_{0 \leq i \leq n_0-1}$ is the column vector whose i th entry is

$$c_i = \sum_{k \geq 0} e_{k-i}^{(m_0+i)}(\mathbf{y}^{(m_0+i-1)}, Y_{m_0+i}).$$

Proof of Theorem 7.6. Let $d \in \mathcal{D}_{n,m}^R$. By the bijection $\Phi : d \mapsto (c_0, c_1)$ in Theorem 7.2 and the fact that $\mathbf{t}^{\bar{U}(d)} \mathbf{x}^d = \mathbf{t}^{\bar{U}(c_0)} \mathbf{x}^{c_0} \cdot \mathbf{t}^{\bar{U}(c_1)} \mathbf{x}^{c_1}$, we obtain that $\sum_{d \in \mathcal{D}_{n,m}^R} \mathbf{t}^{\bar{U}(d)} \mathbf{x}^d = \sum_{(c_0, c_1) \in \mathcal{D}_{n,m}^H} \mathbf{t}^{\bar{U}(c_0)} \mathbf{x}^{c_0} \cdot \mathbf{t}^{\bar{U}(c_1)} \mathbf{x}^{c_1}$. Write $N_0 = n_0 + m_0$ and $N_1 = n_1 + m_1$ for short. Let $\lambda^{(0)} = \text{sh}(c_0)'$ and $\lambda^{(1)} = \text{sh}(c_1)'$. Note that $\text{sh}(c_1) \setminus \text{sh}(c_0)$ is a horizontal strip

if, and only if, $\lambda^{(1)} \setminus \lambda^{(0)}$ is a vertical strip. From Lemma 7.5 above, we conclude that $\sum_{d \in \mathcal{D}_{n,m}^{\mathbb{R}}} \mathbf{t}^{\bar{U}(d)} \mathbf{x}^d$ is equal to

$$\sum_{1 \leq i, j \leq n_0} \det \left(e_{\lambda_j^{(0)} - j + i}^{(N_0 - i)}(\mathbf{y}^{(N_0 - i - 1)}, Y_{N_0 - i}) \right)_{1 \leq i, j \leq n_1} \det \left(e_{\lambda_j^{(1)} - j + i}^{(N_1 - i)}(\mathbf{y}^{(N_1 - i - 1)}, Y_{N_1 - i}) \right),$$

where the sum runs over all pair $(\lambda^{(0)}, \lambda^{(1)})$ of partitions such that $\lambda^{(0)} \subseteq \lambda^{(1)}$ and $\lambda^{(1)} \setminus \lambda^{(0)}$ is a vertical strip. Rearrange the row and column indices, then this equals

$$\sum_{0 \leq i, j \leq n_0 - 1} \det \left(e_{\lambda_{n_0 - j}^{(0)} + j - i}^{(m_0 + i)}(\mathbf{y}^{(m_0 + i - 1)}, Y_{m_0 + i}) \right)_{0 \leq i, j \leq n_1 - 1} \det \left(e_{\lambda_{n_1 - j}^{(1)} + j - i}^{(m_1 + i)}(\mathbf{y}^{(m_1 + i - 1)}, Y_{m_1 + i}) \right).$$

If $n_0 = n_1$, put $l_{j+1}^{(0)} = \lambda_{n_0 - j}^{(0)} + j$ and $l_{j+1}^{(1)} = \lambda_{n_1 - j}^{(1)} + j$ for $0 \leq j \leq n_0 - 1$. Note that $0 \leq l_1^{(k)} < \dots < l_{n_0}^{(k)}$ for $k = 0, 1$. It is easy to see that $\lambda^{(1)} \setminus \lambda^{(0)}$ is a vertical strip if, and only if, $l_j^{(1)} - 1 \leq l_j^{(0)} \leq l_j^{(1)}$. Thus, if we put

$$P_{ij} = e_{j-i}^{(m_1 + i)}(\mathbf{y}^{(m_1 + i - 1)}, Y_{m_1 + i})$$

and

$$\begin{aligned} Q_{ij} &= e_{j-i}^{(m_0 + i)}(\mathbf{y}^{(m_0 + i - 1)}, Y_{m_0 + i}) + e_{j-i-1}^{(m_0 + i)}(\mathbf{y}^{(m_0 + i - 1)}, Y_{m_0 + i}) \\ &= e_{j-i}^{(m_0 + i + 1)}(\mathbf{y}^{(m_0 + i - 1)}, Y_{m_0 + i}, 1) \end{aligned}$$

then, by Lemma 7.4(ii), we obtain $\sum_{d \in \mathcal{D}_{n,m}^{\mathbb{R}}} \mathbf{t}^{\bar{U}(d)} \mathbf{x}^d = \det(Q_{ij})^t (P_{ij})$, which leads to the desired identity (7.7). If $n_0 = n_1 + 1$, then use

$$\det(A) = \det \left(\begin{array}{c|c} 1 & O \\ \hline O & A \end{array} \right)$$

and repeat the same arguments to obtain (7.8). The other identities (7.5) (7.6) can be derived similarly using Lemma 7.4(i). \square

If we specialize the variables in Theorem 7.6, then we obtain the following corollary:

Corollary 7.7. Let m and $n \geq 1$ be nonnegative integers, and fix a positive integer $k \geq 1$. Put $n_0 = \lfloor \frac{n}{2} \rfloor$, $m_0 = \lfloor \frac{n+m+1}{2} \rfloor - n_0$, $n_1 = \lfloor \frac{n}{2} \rfloor$ and $m_1 = \lfloor \frac{n+m+1}{2} \rfloor - n_1$.

- (i) If n is even (i.e. $n_0 = n_1$), then we obtain $\sum_{d \in \mathcal{D}_{n,m}^{\mathbb{R}}} t^{\bar{U}_k(d)} = \det R'(t)$, where $R'(t) = (R'_{ij})_{0 \leq i \leq n_0 - 1, 0 \leq j \leq n_1 - 1}$ is the $n_0 \times n_1$ matrix whose (i, j) th entry is

$$\begin{aligned} R'_{ij} &= \begin{pmatrix} m_0 + m_1 + i + j - 1 \\ m_0 + 2i - j \end{pmatrix} (1 + t^2) \\ &\quad + \left\{ \begin{pmatrix} m_0 + m_1 + i + j - 1 \\ m_0 + 2i - j - 1 \end{pmatrix} + \begin{pmatrix} m_0 + m_1 + i + j - 1 \\ m_0 + 2i - j + 1 \end{pmatrix} \right\} t \end{aligned} \quad (7.9)$$

if $m_0 + i > 0$ and $m_1 + j > 0$, $R'_{0,j} = \begin{pmatrix} m_1 + j \\ -j + 1 \end{pmatrix} + \begin{pmatrix} m_1 + j \\ -j \end{pmatrix} t$ if $m_0 = 0$ and $m_1 + j > 0$, $R'_{i,0} = \delta_{0,i}$ if and $m_0 + i \geq 0$ and $m_1 = 0$.

- (ii) If n is odd (i.e. $n_0 = n_1 + 1$), then we obtain $\sum_{d \in \mathcal{D}_{n,m}^{\mathbb{R}}} t^{\bar{U}_k(d)} = \det (\vec{r} \mid R'(t))$, where $R'(t) = (R'_{ij})_{0 \leq i \leq n_0-1, 0 \leq j \leq n_1-1}$ is the $n_0 \times n_1$ matrix whose (i, j) th entry is

$$R'_{ij} = \binom{m_0 + m_1 + i + j - 1}{m_0 + 2i - j - 1} (1 + t^2) + \left\{ \binom{m_0 + m_1 + i + j - 1}{m_0 + 2i - j - 2} + \binom{m_0 + m_1 + i + j - 1}{m_0 + 2i - j} \right\} t \quad (7.10)$$

if $m_0 + i > 0$ and $m_1 + j > 0$, $R'_{i,0} = \binom{m_0+i}{-i+1} + \binom{m_0+i}{-i} t$ if $m_0 + i > 0$ and $m_1 = 0$, and $R'_{0,j} = \delta_{0,j}$ if $m_0 = 0$ and $m_1 + j \geq 0$. Here $\vec{r} = (\delta_{i,0})_{0 \leq i \leq n_0-1}$ is as in Theorem 7.6.

- (iii) If n is even (i.e. $n_0 = n_1$), then we obtain $\sum_{d \in \mathcal{D}_{n,m}^{\mathbb{C}}} t^{\bar{U}_k(d)} = \det C'(t)$, where $C'(t) = (C'_{ij})_{0 \leq i \leq n_0-1, 0 \leq j \leq n_1-1}$ is the $n_0 \times n_1$ matrix whose (i, j) th entry is

$$C'_{ij} = \sum_{k \leq m_0 + 2i - j - 1} \left[\binom{m_0 + m_1 + i + j - 2}{k} (1 + t^2) + \left\{ \binom{m_0 + m_1 + i + j - 2}{k - 1} + \binom{m_0 + m_1 + i + j - 2}{k + 1} \right\} t \right] \quad (7.11)$$

if $m_0 + i > 0$ and $m_1 + j > 0$, $C'_{i,0} = 2^{m_0+i-1}(1+t)$ if $m_0 + i > 0$ and $m_1 = 0$ and $C'_{0,j} = \delta_{0,j}$ if $m_0 = 0$ and $m_1 + j \geq 0$.

- (iv) If n is odd (i.e. $n_0 = n_1 + 1$), then we obtain $\sum_{d \in \mathcal{D}_{n,m}^{\mathbb{C}}} t^{\bar{U}_k(d)} = \det (\vec{c}'(t) \mid C'(t))$, where $C'(t) = (C'_{ij})_{0 \leq i \leq n_0-1, 0 \leq j \leq n_1-1}$ is the $n_0 \times n_1$ matrix whose (i, j) th entry is

$$C'_{ij} = \sum_{k < m_0 + 2i - j - 1} \left[\binom{m_0 + m_1 + i + j - 2}{k} (1 + t^2) + \left\{ \binom{m_0 + m_1 + i + j - 2}{k - 1} + \binom{m_0 + m_1 + i + j - 2}{k + 1} \right\} t \right] \quad (7.12)$$

if $m_0 + i > 0$ and $m_1 + j > 0$, $2^{m_0+i-1}(1+t) - \delta_{i,0}$ if $m_0 + i > 0$ and $m_1 + j = 0$, and 0 if $m_0 + i = 0$ and $m_1 + j \geq 0$. Here $\vec{c}'(t) = (c'_i)_{0 \leq i \leq n_0-1}$ is the column

vector whose i th entry is $c'_i = \begin{cases} 1, & \text{if } m_0 = i = 0, \\ 2^{m_0+i-1}(1+t), & \text{otherwise.} \end{cases}$

For example, if $n = 5$ and $m = 0$, then $n_0 = 3$, $m_0 = 0$, $n_1 = 2$, $m_1 = 1$, and $\sum_{d \in \mathcal{D}_5^{\mathbb{R}}} t^{\bar{U}_k(d)}$, $k \geq 1$, is equal to

$$\det \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1+t+t^2 & 1+2t+t^2 \\ 0 & t & 3+4t+3t^2 \end{pmatrix} = 3 + 6t + 8t^2 + 6t^3 + 3t^4,$$

whereas $\sum_{d \in \mathcal{D}_5^{\mathbb{C}}} t^{\bar{U}_k(d)}$, $k \geq 1$, is equal to

$$\det \begin{pmatrix} 1 & 0 & 0 \\ 1+t & 1+t+t^2 & t \\ 2+2t & 2+4t+2t^2 & 3+5t+3t^2 \end{pmatrix} = 3 + 6t + 7t^2 + 6t^3 + 3t^4.$$

Proof of Corollary 7.7. Substitute $x_i = 1$ for $i \geq 1$, $t_k = t$ and $t_i = 1$ for $i \neq k$

into Theorem 7.6. First assume $n_0 = n_1$. Using $e_r^n(t, 1, \dots, 1) = \begin{cases} \delta_{0,r}, & \text{if } n = 0, \\ \binom{n-1}{r} + \binom{n-1}{r-1}t, & \text{if } n > 0, \end{cases}$

we obtain

$$R'_{i,j} = \sum_{k \geq 0} \left\{ \binom{m_0 + i}{k - i} + \binom{m_0 + i}{k - i - 1} t \right\} \left\{ \binom{m_1 + j - 1}{k - j} + \binom{m_1 + j - 1}{k - i - 1} t \right\}$$

if $m_1 + j > 0$. Use the binomial coefficient identity $\sum_{k \geq 0} \binom{\alpha}{k} \binom{\beta}{\gamma - k} = \binom{\alpha + \beta}{\gamma}$ to obtain (7.9). The case of $m_1 + j = 0$ is treated similarly. One can prove (7.10) similarly in the case of $n_0 = n_1 + 1$. Next, to prove (ii)(a), substitute $x_i = 1$ for $i \geq 1$, $t_k = t$ and $t_i = 1$ for $i \neq k$ into (7.7). If $m_0 + i > 0$ and $m_1 + j > 0$, then we obtain

$$C'_{i,j} = \sum_{k \geq 0} \sum_{\nu \geq 0} \left\{ \binom{m_0 + i - 1}{k - i} + \binom{m_0 + i - 1}{k - i - 1} t \right\} \left\{ \binom{m_1 + j - 1}{\nu - j} + \binom{m_1 + j - 1}{\nu - i - 1} t \right\}$$

Thus, the proof of the desired identity (7.11) reduce to the following identity

$$\sum_{k \geq 0} \sum_{\nu \geq 0} \binom{m_0 + i - 1}{k - i - \alpha} \binom{m_1 + j - 1}{\nu - j - \beta} = \sum_{k < m_0 + 2i - j + \alpha - \beta} \binom{m_0 + m_1 + i + j - 2}{k}.$$

The other cases can be treated similarly. \square

If we put $m = 0$ in Corollary 7.7, then we obtain the following corollary:

Corollary 7.8. Let n be a positive integer, and fix a positive integer $k \geq 1$. Put $n_0 = \lfloor \frac{n}{2} \rfloor$.

- (i) If n is even, then we obtain $\sum_{d \in \mathcal{Q}_n^R} t^{\bar{U}_k(d)} = \det R_{n_0}^e(t)$, where $R_{n_0}^e(t) = (R_{ij}^e)_{1 \leq i, j \leq n_0 - 1}$ is the $n_0 \times n_0$ matrix whose (i, j) th entry is

$$R_{ij}^e = \binom{i + j}{2i - j + 1} (1 + t^2) + \left\{ \binom{i + j}{2i - j} + \binom{i + j}{2i - j + 2} \right\} t \quad (7.13)$$

if $(i, j) \neq (0, 0)$ and $R_{0,0}^e = 1$.

- (ii) If n is odd, then we obtain $\sum_{d \in \mathcal{Q}_n^R} t^{\bar{U}_k(d)} = \det R_{n_0}^o(t)$, where $R_{n_0}^o(t) = (R_{ij}^o)_{0 \leq i, j \leq n_0 - 1}$ is the $n_0 \times n_0$ matrix whose (i, j) th entry is

$$R_{ij}^o = \binom{i + j - 1}{2i - j} (1 + t^2) + \left\{ \binom{i + j - 1}{2i - j - 1} + \binom{i + j - 1}{2i - j + 1} \right\} t \quad (7.14)$$

if $(i, j) \neq (0, 0), (0, 1)$ and $R_{0,0}^o = R_{0,1}^o = 1$.

- (iii) If n is even, then we obtain $\sum_{d \in \mathcal{Q}_n^C} t^{\bar{U}_k(d)} = \det C_{n_0}^e(t)$, where $C_{n_0}^e(t) = (C_{ij}^e)_{0 \leq i, j \leq n_0 - 1}$ is the $n_0 \times n_0$ matrix whose (i, j) th entry is

$$C_{ij}^e = \left\{ 2 \binom{i + j - 2}{2i - j - 1} + \binom{i + j - 2}{2i - j} \right\} (1 + t^2) \quad (7.15)$$

$$+ \left\{ 2 \binom{i + j - 2}{2i - j - 2} + \binom{i + j - 2}{2i - j - 1} + 2 \binom{i + j - 2}{2i - j} + \binom{i + j - 2}{2i - j + 1} \right\} t$$

if $i + j \geq 2$, $C_{0,0}^e = 1 + t$, $C_{0,1}^e = t$ and $C_{1,0}^e = 0$.

(iv) If n is odd, then we obtain $\sum_{d \in \mathcal{D}_n^C} t^{\bar{U}_k(d)} = \det C_{n_0}^o(t)$, where $C_{n_0}^o(t) = (C_{ij}^o)_{0 \leq i, j \leq n_0-1}$ is the $n_0 \times n_0$ matrix whose (i, j) th entry is

$$C_{ij}^o = \left\{ 2 \binom{i+j-3}{2i-j-2} + \binom{i+j-3}{2i-j-1} \right\} (1+t^2) + \left\{ 2 \binom{i+j-3}{2i-j-3} + \binom{i+j-3}{2i-j-2} + 2 \binom{i+j-3}{2i-j-1} + \binom{i+j-3}{2i-j} \right\} t \quad (7.16)$$

if $i+j \geq 3$, $C_{0,0}^o = 1$, $C_{0,1}^o = C_{0,2}^o = C_{2,0}^o = 0$, $C_{1,0}^o = 1+t$ and $C_{1,1}^o = 1+t+t^2$.

Proof. Substitute $m = 0$ in Corollary 7.7. Then we have $m_0 = 1$ and $m_1 = 0$ if n is even, and we have $m_0 = 0$ and $m_1 = 1$ if n is odd. Using this, we can directly compute each entry of $R_{n_0}^e(t) = R'(t)$ (resp. $R_{n_0}^o(t) = (\vec{r} | R'(t))$) in (i)(ii) of Corollary 7.7 if n is even (resp. odd). We define $n \times n$ matrices U_n^e and U_n^o by $U_n^e = (\delta_{i,j} - 2\delta_{i+1,j})_{0 \leq i, j \leq n-1}$ and $U_n^o = U_n^e + (2\delta_{i^2+(j-1)^2, 0})_{0 \leq i, j \leq n-1}$. As before we substitute $m = 0$ in (iii)(iv) of Corollary 7.7. Then put $C_{n_0}^e(t) = U_{n_0}^e C'(t)$ (resp. $C_{n_0}^o(t) = U_{n_0}^o (\vec{r} | R'(t))$) if n is even (resp. odd). We can compute each entry of $C_{n_0}^e(t)$ and $C_{n_0}^o(t)$ directly. \square

For instance, in the case where $n = 5$ and $n = 6$, R_3^o and R_3^e are

$$\begin{pmatrix} 1 & 1 & 0 \\ 0 & 1+t+t^2 & 1+2t+t^2 \\ 0 & t & 3+4t+3t^2 \end{pmatrix}, \quad \begin{pmatrix} 1 & 1+t+t^2 & t \\ 0 & 1+2t+t^2 & 3+4t+3t^2 \\ 0 & t & 4+7t+4t^2 \end{pmatrix},$$

and C^o and C^e are as follows:

$$\begin{pmatrix} 1 & 0 & 0 \\ 1+t & 1+t+t^2 & t \\ 0 & 2t & 3+3t+3t^2 \end{pmatrix}, \quad \begin{pmatrix} 1+t & t & 0 \\ 0 & 2+t+2t^2 & 1+3t+t^2 \\ 0 & 2t & 5+6t+5t^2 \end{pmatrix}.$$

Thus we obtain certain determinantal formulae for the generating functions of \mathcal{D}_n^R and \mathcal{D}_n^C weighted by \bar{U}_k . Thus the above (ii) with Theorem 3.3, Theorem 6.2 and Theorem 7.2 proves Theorem 1.3. Here the problem of evaluations still remains:

Conjecture 7.9. Let r be a positive integer. Let R_r^o , C_r^e and C_r^o as in Corollary 7.8. Then the following identities would hold.

(i) For $r \geq 1$,

$$\det R_r^o(t) = A_{2r+1}^{\text{VS}}(t). \quad (7.17)$$

(ii) For $r \geq 1$,

$$\det C_r^e(t) = A_{2r}^{\text{HTS}}(t). \quad (7.18)$$

(iii) For $r \geq 1$,

$$\det C_r^o(t) = A_{2r-1}^{\text{HTS}}(t). \quad (7.19)$$

By Theorem 3.3 and Theorem 6.2, Conjecture 1.2 (Conjecture 6 of [17]) reduce to prove the determinantal formula in (7.17) of Conjecture 7.9. We can prove all these identities in a weak form where $t = 1$ by reducing the determinants to the Andrews-Burge Theorem (Lemma 7.10). Here we use the notation $(A)_j = A(A+1) \cdots (A+j-1)$.

Lemma 7.10. (Andrews-Burge [4]) Let

$$M_n(x, y) = \det \left(\binom{i+j+x}{2i-j} + \binom{i+j+y}{2i-j} \right)_{0 \leq i, j \leq n-1}. \quad (7.20)$$

Then

$$M_n(x, y) = \prod_{k=0}^{n-1} \Delta_{2k}(x+y), \quad (7.21)$$

where $\Delta_0(u) = 2$ and for $j > 0$

$$\Delta_{2j}(u) = \frac{(u+2j+2)_j (\frac{1}{2}u+2j+\frac{3}{2})_{j-1}}{(j)_j (\frac{1}{2}u+j+\frac{3}{2})_{j-1}}. \quad (7.22)$$

Especially, when $y = x$, we obtain the identity

$$m_n(x) = \det \left(\binom{i+j+x}{2i-j} \right)_{0 \leq i, j \leq n-1} = \frac{1}{2^n} \prod_{k=0}^{n-1} \Delta_{2k}(2x) \quad (7.23)$$

(see [1, 2, 3, 18]).

Theorem 7.11. Let n be a positive integer, and put $r = n_0 = \lceil \frac{n}{2} \rceil$.

- (i) The number of elements of \mathcal{D}_n^R is equal to A_{2r+1}^{VS} if $n = 2r - 1$ is odd, i.e. $\det R_r^o(1) = A_{2r+1}^{\text{VS}}$.
- (ii) The number of elements of \mathcal{D}_n^R is equal to $\frac{(3r+2)!(2r+1)!(2r)!}{(4r+2)!(r+1)!(r!)^2} A_{2r+1}^{\text{VS}}$ if $n = 2r$ is even, i.e. $\det R_r^e(1) = \frac{(3r+2)!(2r+1)!(2r)!}{(4r+2)!(r+1)!(r!)^2} A_{2r+1}^{\text{VS}}$.
- (iii) The number of elements of \mathcal{D}_n^C is equal to A_n^{HTS} , i.e. $\det C_r^o(1) = A_{2r-1}^{\text{HTS}}$ and $\det C_r^e(1) = A_{2r}^{\text{HTS}}$.

Proof. If we put $t = 1$ into $R_r^o(t)$ and $R_r^e(t)$, then we obtain

$$\det R_r^o(1) = \det_{0 \leq i, j \leq r-1} \left(\binom{i+j+1}{2i-j+1} \right) = \frac{1}{2^{r-1}} \prod_{k=1}^{r-1} \frac{(2k+4)_k (2k+\frac{5}{2})_{k-1}}{(k)_k (k+\frac{5}{2})_{k-1}},$$

and

$$\det R_r^e(1) = \det_{0 \leq i, j \leq r-1} \left(\binom{i+j+2}{2i-j+2} \right) = \frac{1}{2^{r-1}} \prod_{k=1}^{r-1} \frac{(2k+6)_k (2k+\frac{7}{2})_{k-1}}{(k)_k (k+\frac{7}{2})_{k-1}}$$

from (7.23). A direct computation shows that these products are equal to A_{2r+1}^{VS} and $\frac{(3r+2)!(2r+1)!(2r)!}{(4r+2)!(r+1)!(r!)^2} A_{2r+1}^{\text{VS}}$ respectively. Similarly, if we put $t = 1$ into $C_r^o(t)$ and $C_r^e(t)$, then we obtain

$$\det C_r^o(1) = \det_{0 \leq i, j \leq r-1} \left(\binom{i+j}{2i-j} + \binom{i+j-1}{2i-j-1} \right) = \prod_{k=1}^{r-1} \frac{(2k+1)_k (2k+1)_{k-1}}{(k)_k (k+1)_{k-1}},$$

and

$$\det C_r^e(1) = \det_{0 \leq i, j \leq r-1} \left(\binom{i+j}{2i-j} + \binom{i+j+1}{2i-j+1} \right) = 2 \prod_{k=1}^{r-1} \frac{(2k+3)_k (2k+2)_{k-1}}{(k)_k (k+2)_{k-1}},$$

from (7.21). A direct computation shows that these products are equal to A_{2r-1}^{HTS} and A_{2r}^{HTS} respectively. \square

Thus Theorem 7.11(i) proves Conjecture 6 of [17] is true (i.e. Conjecture 1.2) when $t = 1$.

References

- [1] G.E. Andrews, “Plane partitions (III): the weak Macdonald conjecture”, *Invent. Math.* **53** (1979), 193–225.
- [2] G.E. Andrews, “Plane partitions V: the TSSCPP conjecture”, *J. Combin. Theory Ser. A* **66** (1994), 28–39.
- [3] G.E. Andrews, “Pfaff’s method (I): the Mills-Robbins-Rumsey determinant”, *Discrete Math.* **193** (1998), 43–60.
- [4] G.E. Andrews and W.H. Burge, “Determinant identities”, *Pacific J. Math.* **158** (1993), 1–14.
- [5] E.A. Bender and D.E. Knuth, “Enumeration of plane partitions”, *J. Combin. Theory, Ser. A* **13** (1972), 40–54.
- [6] D.M. Bressoud, *Proofs and Confirmations*, Cambridge U.P.
- [7] C. Carré and B. Leclerc, “Splitting the Square of a Schur Function into its Symmetric and Antisymmetric Parts”, *J. Algebraic Combin.* **4** (1995), 201 – 231.
- [8] M. Ishikawa, “On refined enumerations of totally symmetric self-complementary plane partitions I”, [arXiv:math.CO/0602068](https://arxiv.org/abs/math/0602068).
- [9] M. Ishikawa and M. Wakayama, “Minor summation formula of Pfaffians”, *Linear and Multilinear algebra* **39** (1995), 285–305.
- [10] M. Ishikawa and M. Wakayama, “Applications of the minor summation formula III: Plücker relations, lattice paths and Pfaffians”, [arXiv:math.CO/0312358](https://arxiv.org/abs/math/0312358), *J. Combin. Theory Ser. A.* **113** (2006) 113-155.
- [11] C. Krattenthaler, “Determinant identities and a generalization of the number of totally symmetric self-complementary plane partitions”, *Electron. J. Combin.* **4(1)** (1997), #R27.
- [12] C. Krattenthaler, “Advanced determinant calculus”, *Sem. Lothar. Combin.* **42** (“The Andrews Festschrift”) (1999), Article B42q.
- [13] G. Kuperberg, “Another proof of the alternating-sign matrix conjecture”, *Int. Math. Res. Not.* **3** (1996), 139–150. [arXiv:math.CO/9810091](https://arxiv.org/abs/math/9810091).
- [14] G. Kuperberg, “Symmetry classes of alternating-sign matrices under one roof”, *Ann. of Math.* (2) **156** (2002), 835-866, [arXiv:math.CO/0008184](https://arxiv.org/abs/math/0008184).

- [15] I. G. Macdonald, *Symmetric Functions and Hall Polynomials (2nd ed.)*, Oxford Univ. Press, (1995).
- [16] W.H. Mills, D.P. Robbins and H. Rumsey, “Alternating sign matrices and descending plane partitions”, *J. Combin. Theory Ser. A* **34**, (1983), 340–359.
- [17] W.H. Mills, D.P. Robbins and H. Rumsey, “Self-complementary totally symmetric plane partitions”, *J. Combin. Theory Ser. A* **42**, (1986), 277–292.
- [18] W.H. Mills, D.P. Robbins and H. Rumsey, “Enumeration of a symmetry class of plane partitions”, *Discrete Math.* **67**, (1987), 43–55.
- [19] S. Okada, “Enumeration of symmetry classes of alternating sign matrices and characters of classical groups”, [arXiv:math.CO/0308234](https://arxiv.org/abs/math.CO/0308234), *J. Algebraic Combin.* **23** (2006), 43 – 69.
- [20] A.V. Razumov and Yu. G. Stroganov, “On refined enumerations of some symmetry classes of ASMs”, [arXiv:math-ph/0312071](https://arxiv.org/abs/math-ph/0312071).
- [21] A.V. Razumov and Yu. G. Stroganov, “Enumerations of half-turn symmetric alternating-sign matrices of odd order”, [arXiv:math-ph/0504022](https://arxiv.org/abs/math-ph/0504022).
- [22] D.P. Robbins, “Symmetry classes of alternating sign matrices”, [arXiv:math.CO/0008045](https://arxiv.org/abs/math.CO/0008045).
- [23] R.P. Stanley, “Symmetries of plane partitions”, *J. Combin. Theory Ser. A* **43**, (1986), 103–113.
- [24] R.P. Stanley, *Enumerative combinatorics, Volume II*, Cambridge University Press, (1999).
- [25] D. Stanton and D. White, “A Schensted algorithm for rim hook tableaux”, *J. Combin. Theory Ser. A* **40** (1985), 211 – 247.
- [26] Yu. G. Stroganov, “Izergin-Korepin determinant reloaded” [arXiv:math.CO/0409072](https://arxiv.org/abs/math.CO/0409072).
- [27] D. Zeilberger, “Proof of the refined alternating sign matrix conjecture”, *New York J. Math.* **2** (1996), 59–68.